

**Neomycin Sulfate and Oxytetracycline Cattle Supplement I
Type B Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated cattle feed.

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate.....10 grams/lb.
Oxytetracycline (from oxytetracycline dihydrate base) equivalent to 10 grams oxytetracycline hydrochloride/lb.

GUARANTEED ANALYSIS

Crude Protein, (Min).....XX.XX%
NPN, (Max)¹..... X.XX%
Crude Fat, (Min)..... X.XX%
Crude Fiber, (Max)..... X.XX%
Calcium, (Min)..... X.XX%
Calcium, (Max)..... X.XX%
Phosphorus, (Min)..... X.XX%
Salt, (Min)¹..... X.XX%
Salt, (Max)¹..... X.XX%
Sodium, (Min)²..... X.XX%
Sodium, (Max)²..... X.XX%
Potassium, (Min)¹..... X.XX%
Vitamin A, (Min)³..... ____IU/LB

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

³Other than precursor of Vitamin A, if added

INGREDIENTS

Ingredients as defined by AAFCO.

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below. For the manufacture of Type C medicated feeds of less than 100g/ton neomycin and oxytetracycline from this Type B feed, it is recommended that the Type B feed be further diluted before mixing the Type C feed. An example of further dilution would be a ratio of 1:10 of Type B medicated feed:unmedicated feed.

Indications for Use	Oxytetracycline and Neomycin Sulfate Amount	Mixing Directions
For the reduction of the incidence of liver abscesses in growing cattle over 400 lbs.	75 mg/head/day	Mix 1.5 lb of this Type B medicated feed with 1998.5 lb of non-medicated feed to make one ton of Type C medicated feed containing 15 g/ton of neomycin and oxytetracycline*
For prevention and treatment of the early stages of shipping fever complex	0.5 to 2.0 g/head/day	Mix 10 to 40 lb of this Type B medicated feed with 1990 to 1960 lb of non-medicated feed to make one ton of Type C medicated feed containing 100 to 400 g/ton of neomycin and oxytetracycline*
For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <i>Escherichia coli</i> susceptible to neomycin in calves, beef and non-lactating dairy cattle.	10mg/lb body weight/day	Mix 100 lb of this Type B medicated feed with 1900 lb of non-medicated feed to make one ton of Type C medicated feed containing 1000 g/ton of neomycin and oxytetracycline*
Residue Warning: At 10 mg/lb dosage, withdraw 5 days before slaughter. Zero-day withdrawal period for lower use levels. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residue. A withdrawal period has not been established for use in preruminating calves. Do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.		

*A 500 lb animal consuming 10 lb of this Type C feed will receive the required amount of oxytetracycline and neomycin sulfate.

Net Weight _____lb. (_____kg.)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000