

N-141301-4-0001

Ractopamine and Monensin Med. Tom Turkey Feed

Type C Med Feed

Page 1 of 2

Ractopamine and Monensin Medicated Tom Turkey Feed Type C Medicated Feed

TOMS: For increased rate of weight gain and improved feed efficiency in finishing tom turkeys fed for the last 14 days prior to slaughter, and for the prevention of coccidiosis in growing turkeys caused by *Eimeria adenoeides*, *E. meleagridis* and *E. gallopavonis*.

ACTIVE DRUG INGREDIENT

Ractopamine Hydrochloride.....	4.6 to 11.8 g/ton*
Monensin, USP.....	54-90 g/ton**

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than.....	_____	%
Sodium ² , not more than.....	_____	%

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed continuously as the sole ration for the last 14 days prior to slaughter.

* Final printed label on formulated Type C medicated feed must bear a single drug concentration. No increased benefit has been shown when ractopamine concentrations in the diet are greater than 4.6 g/ton.

**Final printed label on formulated Type C medicated feed must bear a single drug concentration.