

N-141221-A-0000  
 OptaFLEXx Finishing Cattle Feed  
 Type C Med. Feed - weight gain/feed efficiency  
 page 1

Net Weight lb (kg) on bag or bulk  
 Optaflexx™ Finishing Cattle Feed  
 Type C Medicated Feed

For increased rate of weight gain and improved feed efficiency in cattle fed in confinement for slaughter during the last 28 to 42 days on feed.

**ACTIVE DRUG INGREDIENT**

Ractopamine hydrochloride..... 8.2 to 24.6 g/ton  
 [Specify one drug level]

**GUARANTEED ANALYSIS**

Crude Protein, not less than.....	_____	%
Non-Protein Nitrogen (NPN) <sup>1</sup> , not more than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt <sup>2</sup> , not less than.....	_____	%
Salt <sup>2</sup> , not more than.....	_____	%
Sodium <sup>3</sup> , not less than.....	_____	%
Sodium <sup>3</sup> , not more than.....	_____	%
Potassium, not less than.....	_____	%
Vitamin A <sup>2,4</sup> , not less than.....	_____	I.U./lb

<sup>1</sup>When added.

<sup>2</sup>If added.

<sup>3</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup>Other than precursors of Vitamin A.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**DIRECTIONS FOR USE**

Feed continuously to cattle fed in confinement for slaughter as the sole ration for last 28 to 42 days on feed.

**WARNING:** The active ingredient in Optaflexx, ractopamine hydrochloride, is a beta-adrenergic agonist. Individuals with cardiovascular disease should exercise special caution to avoid exposure. Not for use in humans. Keep out of the reach of children. The Optaflexx 45 formulation (Type A Medicated Article) poses a low dust potential under usual conditions of handling and mixing. When mixing and handling Optaflexx, use protective clothing, impervious gloves, protective eye wear, and a NIOSH-approved dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse eyes thoroughly with water. If irritation persists, seek medical attention. The material safety data sheet contains more detailed occupational safety information. To report adverse effects, access medical information, or obtain additional product information, call 1-800-428-4441.

**CAUTION:** Not for animals intended for breeding.

**MANUFACTURED BY**  
 BLUE BIRD FEED MILL  
 Any town, USA 12345

\*Elanco®, Optaflexx™, and the diagonal color bar are trademarks of Eli Lilly and Company.

14 Mar03