

**Neomycin Sulfate and Oxytetracycline Cattle Ration E  
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD)  
drug to use by or on the order of a licensed veterinarian.**

**INDICATIONS FOR USE**

For treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia (shipping fever complex) caused by *Pasteurella multocida* susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin in calves, beef and non-lactating dairy cattle.

**ACTIVE DRUG INGREDIENTS**

Neomycin Sulfate..... grams/ton\*  
Oxytetracycline (from oxytetracycline dihydrate base) equivalent to \_\_\_\_ grams oxytetracycline hydrochloride/ton.\* See FEEDING DIRECTIONS.

**GUARANTEED ANALYSIS**

Crude Protein, (Min).....	XX.XX%
NPN, (Max) <sup>1</sup> .....	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min).....	X.XX%
Calcium, (Max).....	X.XX%
Phosphorus, (Min).....	X.XX%
Salt, (Min) <sup>1</sup> .....	X.XX%
Salt, (Max) <sup>1</sup> .....	X.XX%
Sodium, (Min) <sup>2</sup> .....	X.XX%
Sodium, (Max) <sup>2</sup> .....	X.XX%
Potassium, (Min) <sup>1</sup> .....	X.XX%
Vitamin A, (Min) <sup>3</sup> .....	____IU/LB

<sup>1</sup>If added

<sup>2</sup>Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

<sup>3</sup>Other than precursor of Vitamin A, if added

**INGREDIENTS**

Ingredients as defined by AAFCO.

**FEEDING DIRECTIONS**

Feed continuously at 10 mg/lb body weight/day for 7-14 days as the sole ration; in feed or milk replacers. Treatment should continue 24 to 48 hours beyond remission of disease symptoms. For example, the table below with the stated drug concentration, body weight and feed consumption will provide cattle with 10 mg/lb body weight/day dosage.

<b>Concentration of Oxytetracycline and Neomycin Sulfate (g/ton) in Type C Feed</b>	<b>Body Weight (lb)</b>	<b>Type C Feed Consumption (lb)</b>
1000	100	2
1000	175	3.5
1000	250	5
1000	350	7
1000	500	10
1000	650	13

**RESIDUE WARNING:** Withdraw 5 days before slaughter at 10 mg/lb dosage. Use of more than one product containing neomycin or failure to withdrawal times may result in illegal drug residues. A withdrawal period has not been established for use in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.

\*The final printed medicated feed label must state a single drug concentration.

Net Weight \_\_\_\_\_ lb. ( \_\_\_\_\_ kg.)  
Manufactured by  
Blue Bird Feed Company  
Blue Bird, MD 00000