

**Neomycin Sulfate and Oxytetracycline Calf Milk Replacer or Starter Feed Ration (Calves Up To 250 lbs) G
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use
by or on the order of a licensed veterinarian.**

INDICATIONS FOR USE

For treatment of bacterial enteritis caused by *Escherichia coli* susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin in calves up to 250 lbs for use in milk replacers or starter feeds.

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate.....grams/ton.*
Oxytetracycline hydrochloride.....grams/ton.*
See FEEDING DIRECTIONS.

GUARANTEED ANALYSIS

Crude Protein, (Min).....XX.XX%
NPN, (Max)¹..... X.XX%
Crude Fat, (Min)..... X.XX%
Crude Fiber, (Max)..... X.XX%
Calcium, (Min)..... X.XX%
Calcium, (Max)..... X.XX%
Phosphorus, (Min)..... X.XX%
Salt, (Min)¹..... X.XX%
Salt, (Max)¹..... X.XX%
Sodium, (Min)²..... X.XX%
Sodium, (Max)²..... X.XX%
Potassium, (Min)¹..... X.XX%
Vitamin A, (Min)³..... ___IU/LB

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

³Other than precursor of Vitamin A, if added

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed at 10 mg/lb body weight/day for 7-14 days as the sole ration; in calf milk replacers or starter feeds. Treatment should continue 24 to 48 hours beyond remission of disease symptoms. For example, the table below with the stated drug concentration, body weight and feed consumption will provide calves with 10 mg/lb body weight/day dosage.

Concentration of Oxytetracycline and Neomycin Sulfate (g/ton) in Type C Feed	Body Weight (lb)	Type C Feed Consumption (lb)
1000	100	2
1000	175	3.5
1000	250	5

RESIDUE WARNING: Withdraw 5 days before slaughter at 10 mg/lb dosage. Use of more than one product containing neomycin or failure to withdrawal times may result in illegal drug residues. A withdrawal period has not been established for use in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle. Do not use in female dairy cattle 20 months or age or older.

*The final printed medicated feed label must state a single drug concentration.

Net Weight _____ lb. (_____ kg.)
Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000