

**Neomycin Sulfate and Oxytetracycline Cattle Ration D
Type C Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS FOR USE

For prevention and treatment of the early stages of shipping fever complex

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate..... grams/ton*
Oxytetracycline (from oxytetracycline dihydrate base) equivalent to ____grams oxytetracycline hydrochloride/ton.* See FEEDING DIRECTIONS.

GUARANTEED ANALYSIS

Crude Protein, (Min).....XX.XX%
NPN, (Max)¹..... X.XX%
Crude Fat, (Min)..... X.XX%
Crude Fiber, (Max)..... X.XX%
Calcium, (Min)..... X.XX%
Calcium, (Max)..... X.XX%
Phosphorus, (Min)..... X.XX%
Salt, (Min)¹..... X.XX%
Salt, (Max)¹..... X.XX%
Sodium, (Min)²..... X.XX%
Sodium, (Max)²..... X.XX%
Potassium, (Min)¹..... X.XX%
Vitamin A, (Min)³..... ____IU/LB

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

³Other than precursor of Vitamin A, if added

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed at 0.5-2.0 g/head/day 3-5 days before and after arrival in feedlots. For example, the table below with the stated drug concentration, body weight and feed consumption will provide cattle with 0.5-2.0 g/head/day dosage.

Concentration of Oxytetracycline and Neomycin Sulfate (g/ton) in Type C Feed	Body Weight (lb)	Type C Feed Consumption (lb)
100 – 400	500	10
80 – 320	625	12.5
66 - 266	750	15

RESIDUE WARNING: A withdrawal period has not been established for use in preruminating calves. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residue. Do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.

*The final printed medicated feed label must state a single drug concentration.

Net Weight ____lb. (____kg.)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000