

**Neomycin Sulfate and Oxytetracycline Breeding Swine Ration C
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD)
drug to use by or on the order of a licensed veterinarian.**

INDICATIONS FOR USE

For control and treatment of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by Leptospira pomona susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate..... grams/ton.*
Oxytetracycline (from oxytetracycline dihydrate base) equivalent to ____grams oxytetracycline hydrochloride/ton.*
See FEEDING DIRECTIONS.

GUARANTEED ANALYSIS

Crude Protein, (Min).....	XX.XX%
Lysine, (Min).....	X.XX%
Methionine, (Min).....	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min).....	X.XX%
Calcium, (Max).....	X.XX%
Phosphorus, (Min).....	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%
Selenium, (Min).....	ppm
Zinc, (Min).....	ppm

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed at 10mg/lb body weight/day continuously for not more than 14 days as the sole ration. For example, the table below with the stated drug concentration, body weight and feed consumption will provide swine with 10 mg/lb body weight/day dosage.

Concentration of Oxytetracycline and Neomycin Sulfate (g/ton) in Type C feed	Body Weight (lb)	Type C Feed Consumption (lb)
400	100	5
500	150	6.00
600	200	6.67

RESIDUE WARNING: Withdraw 5 days before slaughter.

*The final printed medicated feed label must state a single drug concentration.

Net Weight ____lb. (____kg.)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000