

**Neomycin Sulfate and Oxytetracycline Swine Supplement
Type B Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated swine feed.

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate.....10 grams/lb.
Oxytetracycline (from oxytetracycline dihydrate base) equivalent to 10 grams oxytetracycline hydrochloride/lb.

GUARANTEED ANALYSIS

Crude Protein, (Min).....	XX.XX%
Lysine, (Min).....	X.XX%
Methionine, (Min).....	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min).....	X.XX%
Calcium, (Max).....	X.XX%
Phosphorus, (Min).....	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%
Selenium, (Min).....	ppm
Zinc, (Min).....	ppm

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO.

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below.

Indications for Use	Oxytetracycline and Neomycin Sulfate Amount	Mixing Directions
Swine: For treatment of bacterial enteritis caused by <u>Escherichia coli</u> and <u>Salmonella choleraesuis</u> and bacterial pneumonia caused by <u>Pasteurella multocida</u> susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <u>Escherichia coli</u> susceptible to neomycin.	10 mg/lb body weight/day	Mix 40 lb of this Type B medicated feed with 1960 lb of non-medicated feed to make one ton of Type C medicated feed containing 400 g/ton of neomycin and oxytetracycline*
Breeding Swine: For control and treatment of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <u>Leptospira pomona</u> susceptible to oxytetracycline.	10 mg/lb body weight/day	Mix 40 lb of this Type B medicated feed with 1960 lb of non-medicated feed to make one ton of Type C medicated feed containing 400 g/ton of neomycin and oxytetracycline*
Residue Warning: Withdraw 5 days before slaughter.		

*A 100lb animal consuming 5lb of this Type C feed will receive the required amount of oxytetracycline and neomycin sulfate.

Net Weight _____lb. (_____kg.)
Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000