

**Neomycin Sulfate and Oxytetracycline Chicken Ration B  
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD)  
drug to use by or on the order of a licensed veterinarian.**

**INDICATIONS FOR USE**

For control of infectious synovitis caused by Mycoplasma synoviae; control of fowl cholera caused by Pasteurella multocida susceptible to oxytetracycline.

**ACTIVE DRUG INGREDIENTS**

Neomycin Sulfate.....100-200 grams/ton.\*  
Oxytetracycline (from oxytetracycline dihydrate base) equivalent to 100-200 grams  
oxytetracycline hydrochloride/ton.\*

**GUARANTEED ANALYSIS**

Crude Protein, (Min).....	XX.XX%
Lysine, (Min).....	X.XX%
Methionine, (Min).....	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min).....	X.XX%
Calcium, (Max).....	X.XX%
Phosphorus, (Min).....	X.XX%
Salt, (Min) <sup>1</sup> .....	X.XX%
Salt, (Max) <sup>1</sup> .....	X.XX%
Sodium, (Min) <sup>2</sup> .....	X.XX%
Sodium, (Max) <sup>2</sup> .....	X.XX%

<sup>1</sup>If added

<sup>2</sup>Should be guaranteed only when total sodium exceeds that furnished by  
the maximum salt guarantee

**INGREDIENTS**

Ingredients as defined by AAFCO.

**FEEDING DIRECTIONS**

Feed continuously for 7-14 days as the sole ration.

**RESIDUE WARNING:** Do not feed to chickens producing eggs for human consumption.  
Zero-day withdrawal period. In low calcium feeds, withdraw 3 days prior to slaughter.

\*The final printed medicated feed label must state a single drug concentration.

Net Weight \_\_\_\_\_ lb. ( \_\_\_\_\_ kg.)

Manufactured by  
Blue Bird Feed Company  
Blue bird, MD 00000