

N-140865-E-0001 (CVM Retyped) Approved 8/7/2000  
Narasin/Bacitracin Zinc Broiler Chicken Ration  
Type C Medicated Feed

Net weight lb (kg) on bag or bulk  
**Narasin/Bacitracin Zinc Broiler Chicken Ration**  
**Type C Medicated Feed**

For the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*, and for increased rate of weight gain and improved feed efficiency in broiler chickens.

**ACTIVE DRUG INGREDIENT**

Narasin..... 54 to 72 g/ton  
Bacitracin zinc..... 4 to 50g/ton

**GUARANTEED ANALYSIS**

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt <sup>1</sup> , not less than.....	_____	%
Salt <sup>1</sup> , not more than.....	_____	%
Sodium <sup>2</sup> , not less than.....	_____	%
Sodium <sup>2</sup> , not more than.....	_____	%

<sup>1</sup>If added.

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**DIRECTIONS FOR USE**

Feed continuously as the sole ration.

**CAUTION:** Do not allow adult turkeys, horses or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal.

**MANUFACTURED BY**

BLUE BIRD FEED MILL  
Anytown, USA 12345