

N-140926 - E-0001 (CVM Retyped) Approved 1/4/1999

Blue Bird Broiler Ration Type C Med. Feed

(Narasin 27-45 g/ton) Nicarbazine 27-45 g/ton; Bacitracin methelene disalicylate 4-50 g/ton (Cocci/wt/FE)

Lot Number _____

page 1 of 2

**NET WEIGHT ON BAG OR BULK
BLUE BIRD BROILER RATION
TYPE C MEDICATED FEED**

For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, and for increased rate of weight gain and improved feed efficiency in broiler chickens.

ACTIVE DRUG INGREDIENT

Narasin.....	27 to 45 g/ton
Nicarbazin.....	27 to 45 g/ton
Bacitracin methylene disalicylate.....	4 to 50 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____ %
Lysine, not less than.....	_____ %
Methionine, not less than.....	_____ %
Crude Fat, not less than.....	_____ %
Crude Fiber, not more than.....	_____ %
Calcium, not less than.....	_____ %
Calcium, not more than.....	_____ %
Phosphorus, not less than.....	_____ %
Salt ¹ , not less than.....	_____ %
Salt ¹ , not more than.....	_____ %
Sodium ² , not less than.....	_____ %
Sodium ² , not more than.....	_____ %

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials' Publication (AAFCO Official Publication).

N-140926-E-0011 (CVM Retyped) Approved 1/4/1999

Blue Bird Broiler Ration Type C Med. Feed

Narasin 27-45g/ton; Nicarbazin 27-45g/ton; Bacitracin methylene disalicylate 4-50g/ton
(Cocci 1W6/FE)

FEEDING DIRECTIONS

-- page 2 of 2

Feed continuously as the sole ration.

WARNING: Withdraw 5 days before slaughter

CAUTION: For broiler chickens only. Do not feed to laying hens. Do not allow adult turkeys, horses or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal. Nicarbazin medicated broilers may show reduced heat tolerance if exposed to high temperature and high humidity. Provide adequate drinking water and ventilation during these periods.

MANUFACTURED BY

BLUE BIRD FEED MILL

Anytown, USA 12345