

A-200479-A-0000-0T
ZILPATEROL AND MONENSIN
TYPE C MEDICATED FEED

**Zilpaterol and Monensin
Type C Medicated Cattle Feed**

For Use in Cattle Feeds Only

For increased rate of weight gain, improved feed efficiency, increased carcass leanness, and prevention and control of coccidiosis due to *Eimeria bovis* and *E. zuernii* in cattle fed in confinement for slaughter for the last 20 to 40 days on feed.

ACTIVE DRUG INGREDIENTS

Zilpaterol hydrochloride 6.8 g/ton
Monensin, USP 10 to 40 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than %
Non-Protein Nitrogen (NPN)¹, not more than %
Crude Fat, not less than %
Crude Fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Salt², not less than %
Salt², not more than %
Sodium³, not less than %
Sodium³, not more than %
Potassium, not less than %
Vitamin A^{2,4}, not less than I.U./lb.

¹When added.

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed continuously as sole ration to provide 60 to 90 mg/hd/day zilpaterol and 0.14 to 0.42 mg monensin/lb body weight per day, depending on the severity of the coccidiosis challenge, up to 480 mg/hd/day for the last 20 to 40 days on feed.

*Final printed label on formulated Type C medicated feed must bear a single concentration of each drug

CAUTION

Do not allow horses or other equines access to feeds containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Zilpaterol hydrochloride is not for animals intended for breeding. Do not allow horses or other equine access to feed containing zilpaterol.

WARNING

A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

The active ingredient in Zilmax[®] is zilpaterol hydrochloride, a beta₂-adrenergic agonist. Not for use in humans. An anti-dust process has been applied to the drug product, Zilmax[®], in order to greatly reduce inhalation risk. Extended handling tasks with the potential for dust generation require respiratory protection. Wear appropriate skin protection (e.g., impervious gloves, apron, overalls) if there is a potential for extended skin contact. Wear protective eyewear, if there is a potential for eye contact. If accidental eye contact occurs, immediately rinse with water and consult a physician.

→ **WITHDRAWAL PERIOD** 3 days ←

YOU MAY NOTICE: Animals receiving zilpaterol hydrochloride may exhibit increased respiratory rate as well as elevated levels of creatine phosphokinase (CPK) and creatinine

MANUFACTURED BY

BLUE BIRD FEED MILL
Any town, USA 12345

Zilmax[®] is the trademark of Intervet.

Net Weight on Bulk Invoice

01 82 18 00c