

N-141136-M-0022 (CVH retyped)
 Salinomycin/Bacitracin methylene disalicylate (100-200 g/ton) Broiler, Roaster, and Replacement (Breeder and Layer) Chicken Ration
 Type C Med. Feed

Net weight lb (kg) on bag or bulk

**Salinomycin/Bacitracin methylene disalicylate
 Broiler, Roaster, and Replacement (Breeder and Layer) Chicken Ration
 Type C Medicated Feed**

For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, and as an aid in the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to bacitracin in broiler, roaster, and replacement (breeder and layer) chickens.

ACTIVE DRUG INGREDIENTS

Salinomycin.....	40 to 60 g/ton
Bacitracin methylene disalicylate.....	100 to 200 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____ %
Lysine, not less than.....	_____ %
Methionine, not less than.....	_____ %
Crude Fat, not less than.....	_____ %
Crude Fiber, not more than.....	_____ %
Calcium, not less than.....	_____ %
Calcium, not more than.....	_____ %
Phosphorus, not less than.....	_____ %
Salt ¹ , not less than.....	_____ %
Salt ¹ , not more than.....	_____ %
Sodium ² , not less than.....	_____ %
Sodium ² , not more than.....	_____ %

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

DIRECTIONS FOR USE

To control an outbreak of necrotic enteritis, start medication at first clinical signs of disease. The dosage range permitted provides for different levels based on severity of infection. Administer continuously for 5 to 7 days or as long as clinical signs persist, then reduce bacitracin methylene disalicylate to prevention level (50 g/ton). Discontinue use prior to sexual maturity.

CAUTION: For broiler, roaster, and replacement (breeder and layer) chickens only. Do not feed to laying chickens. May be fatal if fed to adult turkeys or to horses.

**MANUFACTURED BY
 BLUE BIRD FEED MILL
 Anytown, USA 12345**