

A-200479- A-0000-OT
HEIFER SUPPLEMENT
TYPE C MEDICATED FEED

**THIS BLUEBIRD LABEL IS INTENDED FOR USE ONLY WITH
MONENSIN AND ZILPATEROL**

BAG OR BULK

**HEIFER SUPPLEMENT
Type C Medicated Feed**

For Beef and Dairy Heifers

INDICATIONS

Heifers Fed in Confinement for Slaughter: For Increased Rate of Weight Gain, Improved Feed Efficiency and Suppression of Estrus (Heat).

ACTIVE DRUG INGREDIENT

Melengestrol acetate0.125 - 1.0 mg/lb.*

GUARANTEED ANALYSIS

| | | |
|---|-------|----------|
| Crude Protein, not less than | _____ | % |
| Non-Protein Nitrogen (NPN) ¹ , not more than | _____ | % |
| Crude Fat, not less than | _____ | % |
| Crude Fiber, not more than | _____ | % |
| Calcium, not less than | _____ | % |
| Calcium, not more than | _____ | % |
| Phosphorus, not less than | _____ | % |
| Salt ² , not less than | _____ | % |
| Salt ² , not more than | _____ | % |
| Sodium ³ , not less than | _____ | % |
| Sodium ³ , not more than | _____ | % |
| Potassium, not less than | _____ | % |
| Vitamin A ^{2,4} , not less than | _____ | I.U./lb. |

¹When added.

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

*Final printed label on formulated Type C medicated feed must bear a single concentration of this drug

DIRECTIONS FOR USE

Must be top dressed or mixed with a complete ration containing monensin (10 to 40 g/ton) and zilpaterol (6.8 g/ton).

Feed at the rate of 0.5-2.0 pound(s) per head per day (specify one level) to provide 0.25-0.5 mg melengestrol acetate per head per day (specify one level). Feed melengestrol acetate in this combination for the final 20 to 40 days on feed.

CAUTION

Melengestrol acetate is not effective in steers and spayed heifers.

Withdrawal periods of three to five days or more should be avoided to prevent the possibility that the heifers may come into estrus (heat) at the time of loading. You should consider feeding an approved medicated feed containing melengestrol acetate during the withdrawal time for this product to prevent heifers from coming into estrus at the time of loading.

MANUFACTURED BY

BLUE BIRD FEED MILL
Any town, USA 12345

Lot Number _____

Net Weight on Bulk Invoice