

N-141124-T-0004 Approved: 1/14/2002

Blue Bird Narasin/Nicarbazin/BMD PNE

Type C Broiler Feed Medicated

Lot No. _____

NET WEIGHT ON BAG OR BULK

**BLUE BIRD NARASIN/NICARBAZIN/BMD PNE
TYPE C BROILER FEED
MEDICATED**

For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*; and as an aid in the prevention of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to bacitracin methylene disalicylate

ACTIVE DRUG INGREDIENTS

Narasin	27-45 g/ton
Nicarbazin	27-45 g/ton
Bacitracin methylene disalicylate	50 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than	_____ %
Lysine, not less than	_____ %
Methionine, not less than	_____ %
Crude Fat, not less than	_____ %
Crude Fiber, not more than	_____ %
Calcium, not less than	_____ %
Calcium, not more than	_____ %
Phosphorus, not less than	_____ %
Salt ¹ , not less than	_____ %
Salt ¹ , not more than	_____ %
Sodium ² , not less than	_____ %
Sodium ² , not more than	_____ %

¹If added

²Shall be guaranteed only when total Sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

DIRECTIONS FOR USE

Feed continuously as the sole ration.

WARNING: Withdraw 5 days before slaughter.

CAUTION: Do not allow adult turkeys, horses or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal. Nicarbazin medicated broilers may show reduced heat tolerance if exposed to high temperature and high humidity. Provide adequate drinking water and ventilation during these periods. For broiler chickens only. Do not feed to laying chickens. Not for use in feeds for breeder chickens.

BLUE BIRD FEED MILL
Any Town, USA 12345