

A-200479- A-0000-OT
 LIQUID HEIFER SUPPLEMENT
 TYPE C MEDICATED FEED

**THIS BLUEBIRD LABEL IS INTENDED FOR USE ONLY WITH
 MONENSIN AND ZILPATEROL**

**LIQUID HEIFER SUPPLEMENT
 Type C Medicated Feed**

For Beef and Dairy Heifers

INDICATIONS

Heifers Fed in Confinement for Slaughter: For Increased Rate of Weight Gain, Improved Feed Efficiency and Suppression of Estrus (Heat).

ACTIVE DRUG INGREDIENT

Melengestrol acetate0.125 – 1.0 mg/lb.*

GUARANTEED ANALYSIS

Crude Protein, not less than	_____	%
Non-Protein Nitrogen (NPN) ¹ , not more than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than	_____	%
Salt ² , not less than	_____	%
Salt ² , not more than	_____	%
Sodium ³ , not less than	_____	%
Sodium ³ , not more than	_____	%
Potassium, not less than	_____	%
Vitamin A ^{2,4} , not less than	_____	I.U./lb.
pH	4.0 to 8.0	

¹When added.
²If added.
³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.
⁴Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

*Final printed label on formulated Type C medicated feed must bear a single concentration of this drug

DIRECTIONS FOR USE

Must be top dressed or mixed with a complete ration containing monensin (10 to 40 g/ton) and zilpaterol (6.8 g/ton).

Feed at the rate of 0.5-2.0 pound(s) per head per day (specify one level) to provide 0.25-0.5 mg melengestrol acetate per head per day (specify one level). Feed melengestrol acetate in this combination for the final 20 to 40 days on feed.

MIXING DIRECTIONS

Mixing directions for liquid Type C feeds stored in recirculation tank systems are: "Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1 percent of the tank contents from the bottom of the tank to the top. Recirculate daily, as directed in this paragraph even when the Type C feed is not used." Mixing directions for liquid Type C feeds stored in mechanical, air or other agitation-type tank systems are: "Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily, as directed in this paragraph, even when the Type C feed is not used."

CAUTION

Melengestrol acetate is not effective in steers and spayed heifers.

Withdrawal periods of three to five days or more should be avoided to prevent the possibility that the heifers may come into estrus (heat) at the time of loading. You should consider feeding an approved medicated feed containing melengestrol acetate during the withdrawal time for this product to prevent heifers from coming into estrus at the time of loading.

MANUFACTURED BY

BLUE BIRD FEED MILL
Any town, USA 12345

Lot Number _____

Net Weight on Bulk Invoice

01 82 18 00c