

N141154-E-0003 Approved 11/3/2000

Lot No. _____

Blue Bird Robenidine/BMD-P
Type C Broiler/Fryer Feed
Medicated

NET WEIGHT ON BAG OR BULK

BLUE BIRD ROBENIDINE/BMD-P TYPE C BROILER/FRYER FEED MEDICATED

As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati* and as an aid in the prevention of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to bacitracin.

ACTIVE DRUG INGREDIENTS

Robenidine hydrochloride	30 g/ton
Bacitracin methylene disalicylate	50 g/ton

GUARANTEED ANALYSIS

Crude protein, not less than	%
Lysine, not less than	%
Methionine, not less than	%
Crude fat, not less than	%
Crude fiber, not more than	%
Calcium, not less than	%
Calcium, not more than	%
Phosphorus, not less than	%
Salt ¹ , not less than	%
Salt ¹ , not more than	%
Sodium ² , not less than	%
Sodium ² , not more than	%

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed continuously as the sole ration.

WARNING: Withdraw 5 days before slaughter. Do not feed to laying hens.

CAUTION: For broiler and fryer chickens only. Type C feed containing robenidine hydrochloride must be fed within 50 days from the date of manufacture.

BLUE BIRD FEED MILLS
Any Town, USA 12345