

N-141124-T-0004 Approved 1/14/2002

Blue Bird Narasin/Nicarbazin/BMD CNE

Type C Broiler Feed Medicated

Lot No. _____

NET WEIGHT ON BAG OR BULK

**BLUE BIRD NARASIN/NICARBAZIN/BMD CNE
TYPE C BROILER FEED
MEDICATED**

For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*; and as an aid in the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to bacitracin methylene disalicylate

ACTIVE DRUG INGREDIENTS

| | |
|--|---------------|
| Narasin | 27-45 g/ton |
| Nicarbazin | 27-45 g/ton |
| Bacitracin methylene disalicylate..... | 100-200 g/ton |

GUARANTEED ANALYSIS

| | |
|--|---------|
| Crude Protein, not less than..... | _____ % |
| Lysine, not less than..... | _____ % |
| Methionine, not less than..... | _____ % |
| Crude Fat, not less than..... | _____ % |
| Crude Fiber, not more than..... | _____ % |
| Calcium, not less than..... | _____ % |
| Calcium, not more than..... | _____ % |
| Phosphorus, not less than..... | _____ % |
| Salt ¹ , not less than..... | _____ % |
| Salt ¹ , not more than..... | _____ % |
| Sodium ² , not less than..... | _____ % |
| Sodium ² , not more than..... | _____ % |

¹If added

²Shall be guaranteed only when total Sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

DIRECTIONS FOR USE

Feed continuously as the sole ration. To control necrotic enteritis, start this feed at first clinical signs of disease; vary bacitracin dosage based on severity of infection; administer continuously for 5-7 days or as long as clinical signs persist, then reduce bacitracin to prevention level (50 g/ton).

WARNING: Withdraw 5 days before slaughter.

CAUTION: Do not allow adult turkeys, horses or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal. Nicarbazin medicated broilers may show reduced heat tolerance if exposed to high temperature and high humidity. Provide adequate drinking water and ventilation during these periods. For broiler chickens only. Do not feed to laying chickens. Not for use in feeds for breeder chickens. Consult a poultry diagnostic laboratory or pathologist to determine the diagnosis and advice regarding the optimal level of drug.

BLUE BIRD FEED MILL
Any Town, USA 12345