

Blue Bird Robenidone/BMD-CNE  
Type C Broiler/Fryer Feed  
Medicated

**NET WEIGHT ON BAG OR BULK**  
**BLUE BIRD ROBENIDINE/BMD-CNE**  
**TYPE C BROILER/FRYER FEED**  
**MEDICATED**

As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati* and as an aid in the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to bacitracin.

**ACTIVE DRUG INGREDIENTS**

Robenidine hydrochloride .....	30 g/ton
Bacitracin methylene disalicylate .....	100-200 g/ton

**GUARANTEED ANALYSIS**

Crude protein, not less than .....	_____	%
Lysine, not less than .....	_____	%
Methionine, not less than .....	_____	%
Crude fat, not less than .....	_____	%
Crude fiber, not more than .....	_____	%
Calcium, not less than .....	_____	%
Calcium, not more than .....	_____	%
Phosphorus, not less than .....	_____	%
Salt <sup>1</sup> , not less than .....	_____	%
Salt <sup>1</sup> , not more than .....	_____	%
Sodium <sup>2</sup> , not less than .....	_____	%
Sodium <sup>2</sup> , not more than .....	_____	%

<sup>1</sup>If added.  
<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Ingredients as defined by AAFCO.

**FEEDING DIRECTIONS**

To control necrotic enteritis, start this feed at first clinical signs of disease; vary bacitracin dosage based on severity of infection; administer continuously for 5-7 days or as long as clinical signs persist, then reduce bacitracin to prevention level (50 g/ton). Feed continuously as the sole ration.

**WARNING:** Withdraw 5 days before slaughter. Do not feed to laying hens.

**CAUTION:** For broiler and fryer chickens only. Type C feed containing robenidine hydrochloride must be fed within 50 days from the date of manufacture.

**BLUE BIRD FEED MILLS**  
Any Town, USA 12345