

NADA 200-483  
Zilpaterol Finishing Cattle Feed  
Type C Medicated Feed.

**Zilpaterol Finishing Cattle Feed  
Type C Medicated Feed**

For increased rate of weight gain, improved feed efficiency and increased carcass leanness in cattle fed in confinement for slaughter during the last 20 to 40 days on feed.

**ACTIVE DRUG INGREDIENTS**

Zilpaterol hydrochloride ..... 6.8 g/ton

**GUARANTEED ANALYSIS**

Crude Protein, not less than .....	_____	%
Non-Protein Nitrogen (NPN) <sup>1</sup> , not more than .....	_____	%
Crude Fat, not less than .....	_____	%
Crude Fiber, not more than .....	_____	%
Calcium, not less than .....	_____	%
Calcium, not more than .....	_____	%
Phosphorus, not less than .....	_____	%
Salt <sup>2</sup> , not less than .....	_____	%
Salt <sup>2</sup> , not more than .....	_____	%
Sodium <sup>3</sup> , not less than .....	_____	%
Sodium <sup>3</sup> , not more than .....	_____	%
Potassium, not less than .....	_____	%
Vitamin A <sup>2,4</sup> , not less than .....	_____	I.U./lb.

<sup>1</sup>When added.

<sup>2</sup>If added.

<sup>3</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup>Other than precursors of Vitamin A.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**FEEDING DIRECTIONS**

Feed continuously to cattle fed in confinement for slaughter as the sole ration for last 20 to 40 days on feed to provide 60 to 90 mg zilpaterol per head per day.

**WARNING:** The active ingredient in Zilmax<sup>®</sup> is zilpaterol hydrochloride, a beta<sub>2</sub>-adrenergic agonist. Not for use in humans. An anti-dust process has been applied to the drug product, Zilmax<sup>®</sup>, in order to greatly reduce inhalation risk. Extended handling tasks with the potential for dust generation require respiratory protection. Wear appropriate skin protection (e.g., impervious gloves, apron, overalls) if there is a potential for extended skin contact. Wear protective eyewear, if there is a potential for eye contact. If accidental eye contact occurs, immediately rinse with water and consult a physician.

→ WITHDRAWAL PERIOD 3 days ←

**CAUTION:** Zilpaterol hydrochloride is not for animals intended for breeding. Do not allow horses or other equines access to feed containing zilpaterol. Do not use in veal calves.

**YOU MAY NOTICE:** Animals receiving zilpaterol hydrochloride may exhibit increased respiratory rate as well as elevated levels of creatine phosphokinase (CPK) and creatinine

**MANUFACTURED BY**  
**BLUE BIRD FEED MILL**  
Any town, USA 12345

Zimax<sup>®</sup> is the trademark of Intervet.

**Net Weight on Bulk Invoice**