

ANADA 200-483-A-0000  
Zilpaterol Finishing Cattle feed  
Type B Medicinal Feed  
Page 1 of 2

**Zilpaterol Finishing Cattle Feed  
Type B Medicated Feed**

**Do Not Feed Undiluted.**

Important: Must be thoroughly mixed into feed before use.

For increased rate of weight gain, improved feed efficiency and increased carcass leanness in cattle fed in confinement for slaughter during the last 20 to 40 days on feed.

**ACTIVE DRUG INGREDIENTS**

Zilpaterol hydrochloride .....68 to 680 g/ton\*

**GUARANTEED ANALYSIS**

Crude Protein, not less than .....	_____	%
Non-Protein Nitrogen (NPN) <sup>1</sup> , not more than .....	_____	%
Crude Fat, not less than .....	_____	%
Crude Fiber, not more than .....	_____	%
Calcium, not less than .....	_____	%
Calcium, not more than .....	_____	%
Phosphorus, not less than .....	_____	%
Salt <sup>2</sup> , not less than .....	_____	%
Salt <sup>2</sup> , not more than .....	_____	%
Sodium <sup>3</sup> , not less than .....	_____	%
Sodium <sup>3</sup> , not more than .....	_____	%
Potassium, not less than .....	_____	%
Vitamin A <sup>2,4</sup> , not less than .....	_____	I.U./lb.

<sup>1</sup>When added.

<sup>2</sup>If added.

<sup>3</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup>Other than precursors of Vitamin A.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

\*Final printed label must bear a single drug concentration.

**MIXING DIRECTIONS:**

Thoroughly mix this Type B Medicated Feed into one ton of complete cattle feed according to the table below to obtain the proper concentration in the Type C Medicated Feed.

Concentration of Zilpaterol in Type B Medicated Feed grams/pound	Pounds Type B Medicated Feed to Add Per Ton of Type C Medicated Feed	Resulting Zilpaterol Concentration in Type C Medicated Feed <sup>a</sup> grams/ton
0.034	200	6.8
0.170	40	6.8
0.340	20	6.8

<sup>a</sup>Based on 90% Dry Matter Basis

**WARNING:** The active ingredient in Zilmax<sup>®</sup> is zilpaterol hydrochloride, a beta<sub>2</sub>-adrenergic agonist. Not for use in humans. An anti-dust process has been applied to the drug product, Zilmax<sup>®</sup>, in order to greatly reduce inhalation risk. Extended handling tasks with the potential for dust generation require respiratory protection. Wear appropriate skin protection (e.g., impervious gloves, apron, overalls) if there is a potential for extended skin contact. Wear protective eyewear, if there is a potential for eye contact. If accidental eye contact occurs, immediately rinse with water and consult a physician.

→ WITHDRAWAL PERIOD 3 days ←

**CAUTION:** Not for use in animals intended for breeding. Do not allow horses or other equines access to feed containing zilpaterol. Do not use in veal calves.

**YOU MAY NOTICE:** Animals receiving zilpaterol hydrochloride may exhibit increased respiratory rate as well as elevated levels of creatine phosphokinase (CPK) and creatinine

**MANUFACTURED BY**

BLUE BIRD FEED MILL  
Any town, USA 12345

Zilmax<sup>®</sup> is the trademark of Intervet.

Net Weight on Bulk Invoice