

**Zilpaterol Finishing Cattle Feed
Liquid Type B Medicated Feed**

Do Not Feed Undiluted.

Important: Must be thoroughly mixed into feed before use.

For increased rate of weight gain, improved feed efficiency and increased carcass leanness in cattle fed in confinement for slaughter during the last 20 to 40 days on feed.

ACTIVE DRUG INGREDIENT

Zilpaterol hydrochloride68 to 680 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than	_____	%
Non-Protein Nitrogen (NPN) ¹ , not more than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than	_____	%
Salt ² , not less than	_____	%
Salt ² , not more than	_____	%
Sodium ³ , not less than	_____	%
Sodium ³ , not more than	_____	%
Potassium, not less than	_____	%
Vitamin A ^{2,4} , not less than	_____	I.U./lb.
Dry Matter, not less than	60%	
Dry Matter, not more than	75%	
pH	3.8 to 7.5	

¹When added.

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

*Final printed label must bear a single drug concentration.

MIXING DIRECTIONS

Thoroughly mix this Type B Medicated Feed into one ton of complete cattle feed according to the table below to obtain the proper concentration in the Type C Medicated Feed

Concentration of Zilpaterol in Type B Medicated Feed grams/pound	Pounds Liquid Type B Medicated Feed to Add to Non-Medicated Feed to Make One Ton of Type C Medicated Feed	Resulting Zilpaterol Concentration in Type C Medicated Feed ^a grams/ton
0.034	200	6.8
0.170	40	6.8
0.340	20	6.8

^aBased on 90% Dry Matter Basis

For liquid feeds stored in recirculating tank systems:

Recirculate immediately prior to use for not less than 10 minutes, moving not less than 1 percent of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used.

For liquid feeds stored in mechanical, air or other agitation-type tanks systems:

Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.

WARNING: The active ingredient in Zilmax[®] is zilpaterol hydrochloride, a beta₂-adrenergic agonist. Not for use in humans. An anti-dust process has been applied to the drug product, Zilmax[®], in order to greatly reduce inhalation risk. Extended handling tasks with the potential for dust generation require respiratory protection. Wear appropriate skin protection (e.g., impervious gloves, apron, overalls) if there is a potential for extended skin contact. Wear protective eyewear, if there is a potential for eye contact. If accidental eye contact occurs, immediately rinse with water and consult a physician.

→ WITHDRAWAL PERIOD 3 days ←

CAUTION: Not for use in animals intended for breeding. Do not allow horses or other equines access to feed containing zilpaterol. Do not use in veal calves.

YOU MAY NOTICE: Animals receiving zilpaterol hydrochloride may exhibit increased respiratory rate as well as elevated levels of creatine phosphokinase (CPK) and creatinine.

MANUFACTURED BY

BLUE BIRD FEED MILL
Any town, USA 12345

Zilmax[®] is the trademark of Intervet.

Net Weight on Bulk Invoice

01 84 18 00b