

**THIS BLUEBIRD LABEL IS INTENDED FOR USE ONLY WITH  
ZILPATEROL**

**LIQUID HEIFER SUPPLEMENT  
Type C Medicated Feed**

**For Beef and Dairy Heifers**

**INDICATIONS**

Heifers Fed in Confinement for Slaughter: For Increased Rate of Weight Gain, Improved Feed Efficiency and Suppression of Estrus (Heat).

**ACTIVE DRUG INGREDIENT**

Melengestrol acetate .....0.125 – 1.0 mg/lb.\*

**GUARANTEED ANALYSIS**

Crude Protein, not less than .....	_____	%
Non-Protein Nitrogen (NPN) <sup>1</sup> , not more than .....	_____	%
Crude Fat, not less than .....	_____	%
Crude Fiber, not more than .....	_____	%
Calcium, not less than .....	_____	%
Calcium, not more than .....	_____	%
Phosphorus, not less than .....	_____	%
Salt <sup>2</sup> , not less than .....	_____	%
Salt <sup>2</sup> , not more than .....	_____	%
Sodium <sup>3</sup> , not less than .....	_____	%
Sodium <sup>3</sup> , not more than .....	_____	%
Potassium, not less than .....	_____	%
Vitamin A <sup>2,4</sup> , not less than .....	_____	I.U./lb.
pH .....	4.0 to 8.0	

<sup>1</sup>When added.

<sup>2</sup>If added.

<sup>3</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup>Other than precursors of Vitamin A.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**DIRECTIONS FOR USE**

Must be top dressed or mixed with a complete ration containing zilpaterol (6.8 g/ton).

Feed at the rate of 0.5-2.0 pound(s) per head per day (specify one level) to provide 0.25-0.5 mg melengestrol acetate per head per day (specify one level). Feed melengestrol acetate in this combination for the final 20 to 40 days on feed.

\*Final printed label on formulated Type C medicated feed must bear a single concentration of this drug

### MIXING DIRECTIONS

Mixing directions for liquid Type C feeds stored in recirculation tank systems are:  
"Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1 percent of the tank contents from the bottom of the tank to the top. Recirculate daily, as directed in this paragraph even when the Type C feed is not used." Mixing directions for liquid Type C feeds stored in mechanical, air or other agitation-type tank systems are:  
"Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily, as directed in this paragraph, even when the Type C feed is not used."

### CAUTION

Melengestrol acetate is not effective in steers and spayed heifers.

Withdrawal periods of three to five days or more should be avoided to prevent the possibility that the heifers may come into estrus (heat) at the time of loading. You should consider feeding an approved medicated feed containing melengestrol acetate during the withdrawal time for this product to prevent heifers from coming into estrus at the time of loading.

### MANUFACTURED BY

BLUE BIRD FEED MILL  
Any town, USA 12345

Lot Number \_\_\_\_\_

Net Weight on Bulk Invoice