

N-141146-A-0000 (Cum Retyped) Approved Feb 2, 2000

Nicarbazin/Bacitracin zinc Broiler Chicken Ration

Type C medicated feed

Net Weight lb (kg) on bag or bulk

p182

Nicarbazin/Bacitracin zinc Broiler Chicken Ration Type C Medicated Feed

As an aid in preventing outbreaks of cecal (*Eimeria tenella*) and intestinal (*E. acervulina*, *E. maxima*, *E. necatrix*, and *E. brunetti*) coccidiosis, and for increased rate of weight gain and improved feed efficiency in broiler chickens.

ACTIVE DRUG INGREDIENT

Nicarbazin.....	113.5 g/ton
Bacitracin zinc.....	4 to 50 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than.....	_____	%
Sodium ² , not more than.....	_____	%

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

DIRECTIONS FOR USE

Feed continuously as the sole ration from time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard.

(OVER)

N 141146-A-0000 (CUM Retyped) Approved Feb 2, 2000
Niacarbazin/Bacitracin zinc Broiler Chicken ~~Feed~~ Ration
Type C medicated feed
p 2 of 2

WARNING: Discontinue medication four (4) days before marketing the birds for human consumption to allow for elimination of the drug from the edible tissue.

CAUTION: Do not feed to laying hens in production. Do not use as a treatment for outbreaks of coccidiosis. Broiler chickens fed niacarbazin may show reduced heat tolerance when exposed to high temperature and high humidity to which they have not been accustomed and under severe conditions, fatalities may result. An ample supply of drinking water and adequate ventilation will improve the birds' tolerance of heat.

MANUFACTURED BY

BLUE BIRD FEED MILL

Any town, USA 12345