

CARBADOX – OXYTETRACYCLINE
TYPE C
BLUE BIRD
SWINE FEED
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* susceptible to oxytetracycline and treatment of bacterial pneumonia caused by *Pasteurella multocida* susceptible to oxytetracycline and increased rate of weight gain and improved feed efficiency.

ACTIVE DRUG INGREDIENT

Carbadox.....10 - 25 g/ton
Oxytetracycline.....400 g/ton*

* The oxytetracycline concentration should be selected based on feeding regime and feed consumption and should be adequate to deliver 10 mg/lb body weight daily.

GUARANTEED ANALYSIS

Crude Protein (Min).....%
Lysine (Min).....%
Crude Fat (Min).....%
Crude Fiber (Max).....%
Calcium (Min).....%
Calcium (Max).....%
Phosphorus (Min).....%
Salt, if added (Min).....%
Salt, if added (Max).....%
Sodium (Min)*.....%
Sodium (Max)*.....%
Selenium (Min).....ppm
Zinc (Min).....ppm

* Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed continuously as the sole ration for 7-14 days

CAUTION: Not for use in pregnant swine or swine intended for breeding purposes. Do not mix in feeds containing bentonite.

WARNING: DO NOT FEED TO SWINE WITHIN 42 DAYS OF SLAUGHTER.

____ lbs (____ kg) NET WEIGHT

BLUE BIRD FEED MILL
Robin, IN 00000