

Ivermectin/lincomycin Combination
in Medicated Feed for Swine
Type C Medicated Feed

BLUEBIRD
TYPE C MEDICATED FEED
FOR WEANED, GROWING & FINISHING PIGS

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the treatment and control of gastrointestinal roundworms (*Ascaris suum*, adults and fourth-stage larvae; *Ascarops strongylina*, adults; *Hyostromylus rubidus*, adult and fourth-stage larvae; *Oesophagostomum* spp., adult and fourth-stage larvae), kidneyworms (*Stephanurus dentatus*, adults and fourth-stage larvae), lungworms (*Metastrongylus* spp., adults), lice (*Haematopinus suis*) and mange mites (*Sarcoptes scabiei* var. *suis*).

Treatment of swine dysentery.

ACTIVE DRUG INGREDIENTS

Ivermectin.....1.8 g/ton
Lincomycin.....100.0 g/ton
(as LINCOMIX® brand of lincomycin hydrochloride agricultural grade)

GUARANTEED ANALYSIS

Protein, not less than _%
Fat, not less than _%
Fiber, not more than _%

INGREDIENTS

Each ingredient must be specifically named (unless stated as such in the guaranteed analysis listing) in accordance with the names and definitions adopted by AAFCO. Collective terms as approved by AAFCO may be used where applicable.

FEEDING DIRECTIONS

To provide the recommended level of 100 g lincomycin/ton and 1.8 g ivermectin/ton (approximately 100 mcg ivermectin per kg of bodyweight per day), feed as the only feed for 7 consecutive days. Continue by feeding a separate feed containing 100 g lincomycin/ton to complete the lincomycin treatment.

Ivermectin/lincomycin Combination
in Medicated Feed for Swine
Type C Medicated Feed

WARNING: **Withdraw** 6 days before slaughter.
Keep this and all drugs out of the reach of children.

CAUTION: Not to be fed to swine that weigh more than 250 lbs.

This product contains IVOMEC® (ivermectin) and has been formulated specifically for use in swine only. This product should not be used for other animal species. Do not permit water runoff from swine production sites to directly enter lakes, streams, or ponds. Do not contaminate water by direct application or by the improper disposal of drug containers.

Occasionally, swine fed lincomycin may within the first two days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within five to eight days without discontinuing the lincomycin treatment.

Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

®IVOMEC is a registered trademark of Merial, Inc.,
Duluth, GA 30096; LINCOMIX is a registered
trademark of [ZOETIS Inc.](http://www.zoetis.com), Kalamazoo, MI 49001

Manufactured by

BLUEBIRD FEED COMPANY
Robin, Indiana 20260

Net Weight on bag