

NH41085-R-0010 Approved 11/30/2001  
Blue Bird Zoalene/BMD-~~LM2C~~ LM2C  
Type C Replacement Chicken Grower Feed Medicated Lot No. \_\_\_\_\_

**NET WEIGHT ON BAG OR BULK**

**BLUE BIRD ZOALENE/BMD- LM2C  
TYPE C REPLACEMENT CHICKEN GROWER FEED  
MEDICATED**

Development of active immunity to coccidiosis and as an aid in the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to bacitracin

**ACTIVE DRUG INGREDIENTS**

Zoalene ..... 36.3 to 75.4 g/ton  
Bacitracin methylene disalicylate ..... 100 to 200 g/ton

**GUARANTEED ANALYSIS**

Crude Protein, not less than .....	_____	%
Lysine, not less than .....	_____	%
Methionine, not less than .....	_____	%
Crude Fat, not less than .....	_____	%
Crude Fiber, not more than .....	_____	%
Calcium, not less than .....	_____	%
Calcium, not more than .....	_____	%
Phosphorus, not less than .....	_____	%
Salt <sup>1</sup> , not less than .....	_____	%
Salt <sup>1</sup> , not more than .....	_____	%
Sodium <sup>2</sup> , not less than .....	_____	%
Sodium <sup>2</sup> , not more than .....	_____	%

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total Sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Ingredients as defined by AAFCO.

**FEEDING DIRECTIONS**

Feed continuously as the sole ration where light to moderate exposure to coccidiosis is expected. To control necrotic enteritis, start this feed at first clinical signs of disease; vary bacitracin dosage based on severity of infection; administer continuously for 5-7 days or as long as clinical signs persist, then reduce bacitracin to prevention level (50 g/ton).

**CAUTION:** Do not feed to birds over 14 weeks of age. Consult a veterinarian or pathologist if losses exceed 0.5% in a two-day period.

Blue Bird Feed Mills  
Robin, IN 12345