

N-141085-R-0010 Approved 11/30/2001  
Blue Bird Zoalene/BMD CNE  
Type C Broiler Chicken Feed Medicated

Lot No. \_\_\_\_\_

**NET WEIGHT ON BAG OR BULK**

**BLUE BIRD ZOALENE/BMD CNE  
TYPE C BROILER CHICKEN FEED  
MEDICATED**

For prevention and control of coccidiosis and as an aid in the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to bacitracin.

**ACTIVE DRUG INGREDIENTS**

Zoalene..... 113.5 g/ton  
Bacitracin methylene disalicylate..... 100-200 g/ton

**GUARANTEED ANALYSIS**

Crude Protein, not less than..... %  
Lysine, not less than..... %  
Methionine, not less than..... %  
Crude Fat, not less than..... %  
Crude Fiber, not more than..... %  
Calcium, not less than..... %  
Calcium, not more than..... %  
Phosphorus, not less than..... %  
Salt<sup>1</sup>, not less than..... %  
Salt<sup>1</sup>, not more than..... %  
Sodium<sup>2</sup>, not less than..... %  
Sodium<sup>2</sup>, not more than..... %

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total Sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Ingredients as defined by AAFCO.

**FEEDING DIRECTIONS**

Feed continuously as the sole ration. To control necrotic enteritis, start this feed at first clinical signs of disease; vary bacitracin dosage based on severity of infection; administer continuously for 5-7 days or as long as clinical signs persist, then reduce bacitracin to prevention level (50 g/ton).

Blue Bird Feed Mills  
Robin, IN 12345