

NADA 141153 - A - 0000 Approved Jan 13, 2000 CVM Retyped

Diclazuril/Bacitracin methylene disalicylate Broiler Chicken Ration
Type C Medicated Feed

Net Weight lb (kg) on bag or bulk

Diclazuril/Bacitracin methylene disalicylate Broiler Chicken Ration Type C Medicated Feed

For the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mitis (mivati)*, and *E. maxima*. Because diclazuril is effective against *E. maxima* later in its life cycle, subclinical intestinal lesions may be present for a short time after infection. Diclazuril was shown in studies to reduce lesion scores and improve performance and health of birds challenged with *E. maxima*. For increased rate of weight gain and improved feed efficiency in broiler chickens.

ACTIVE DRUG INGREDIENT

Diclazuril..... 0.91 g/ton (1 ppm)
Bacitracin methylene disalicylate..... 4 to 50 g/ton

GUARANTEED ANALYSIS

| | | |
|--|-------|---|
| Crude Protein, not less than..... | _____ | % |
| Lysine, not less than..... | _____ | % |
| Methionine, not less than..... | _____ | % |
| Crude Fat, not less than..... | _____ | % |
| Crude Fiber, not more than..... | _____ | % |
| Calcium, not less than..... | _____ | % |
| Calcium, not more than..... | _____ | % |
| Phosphorus, not less than..... | _____ | % |
| Salt ¹ , not less than..... | _____ | % |
| Salt ¹ , not more than..... | _____ | % |
| Sodium ² , not less than..... | _____ | % |
| Sodium ² , not more than..... | _____ | % |

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

DIRECTIONS FOR USE

Feed continuously as the sole ration.

WARNING: Do not use in hens producing eggs for human food.

MANUFACTURED BY

BLUE BIRD FEED MILL
Anytown, USA 12345