

Ivermectin/lincomycin Combination
in Medicated Feed for Swine
Type C Medicated Feed

BLUEBIRD
SWINE TYPE B MEDICATED FEED

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the treatment and control of gastrointestinal roundworms (*Ascaris suum*, adults and fourth-stage larvae; *Ascarops strongylina*, adults; *Hyostrongylus rubidus*, adults and fourth-stage larvae; *Oesophagostomum* spp., adults and fourth-stage larvae), kidneyworms (*Stephanurus dentatus*, adults and fourth-stage larvae), lungworms (*Metastrongylus* spp., adults), lice (*Haematopinus suis*) and mange mites (*Sarcoptes scabiei* var. *suis*).

For reduction in the severity of swine mycoplasmal pneumonia caused by *Mycoplasma hyopneumoniae*.

ACTIVE DRUG INGREDIENTS

Ivermectin.....18.2-364 g/ton
Lincomycin..... 1.0-20 g/lb
(as LINCOMIX® brand of lincomycin hydrochloride agricultural grade)

GUARANTEED ANALYSIS

Crude Protein, not less than _%
Crude Fat, not less than _%
Crude Fiber, not more than _%

INGREDIENTS

Each ingredient must be specifically named (unless stated as such in the guaranteed analysis listing) in accordance with the names and definitions adopted by AAFCO. Collective terms as listed in 21 CFR§501.110 may be used where applicable.

MIXING DIRECTIONS

This Type B Medicated Feed must be mixed with feed ingredients (or unmedicated feed) at the rate of 200 to 10 Pounds per ton to manufacture a Type C Medicated Feed containing 1.8 grams of ivermectin and 200 grams of lincomycin per ton.

IMPORTANT

Must be thoroughly mixed in feeds before use.
Store opened bag in dry place to prevent caking.
Store at room temperature.

Ivermectin/lincomycin Combination
in Medicated Feed for Swine
Type C Medicated Feed

DIRECTIONS FOR USE

Weaned, Growing & Finishing Pigs. Feed the Type C Medicated Feed, containing lincomycin at the recommended level of 200 g/ton and ivermectin at the recommended level of 1.8 g/ton (approximately 100 mcg ivermectin per kg bodyweight per day) as the only feed for 7 consecutive days. A Type C medicated feed containing lincomycin at the recommended level of 200 g/ton should be fed as the sole ration for the duration of the lincomycin 21-day treatment for reduction in the severity of swine mycoplasmal pneumonia.

WARNING: **Withdraw** 6 days before slaughter.
 Keep this and all drugs out of the reach of children.

CAUTION: Not to be fed to swine that weigh more than 250 lbs.

This product contains IVOMEC® (ivermectin) and has been formulated specifically for use in swine only. This product should not be used for other animal species. Do not permit water runoff from swine production sites to directly enter lakes, streams, or ponds. Do not contaminate water by direct application or by the improper disposal of drug containers. Dispose of containers in an approved landfill or by incineration.

Occasionally, swine fed lincomycin may within the first two days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within five to eight days without discontinuing the lincomycin treatment.

Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

®IVOMEC is a registered trademark of Merial, Inc.,
Duluth, GA 30096; LINCOMIX is a registered
trademark of [ZOETIS Inc.](#), Kalamazoo, MI 49001

Manufactured by

BLUEBIRD FEED COMPANY
Robin, Indiana 20260

Net Weight on bag