

# Tilmicosin Medicated Swine Feed

## Type B Medicated Feed

**CAUTION:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

### Do not Feed Undiluted

#### INDICATIONS

For the control of swine respiratory disease associated with *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

#### ACTIVE DRUG INGREDIENT

Tilmicosin (as tilmicosin phosphate)<sup>a</sup> .....18,100 to 36,300 grams per ton\*.

#### GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt <sup>1</sup> , not less than.....	_____	%
Salt <sup>1</sup> , not more than.....	_____	%
Sodium <sup>2</sup> , not less than.....	_____	%
Sodium <sup>2</sup> , not more than.....	_____	%
Selenium, not less than.....	_____	ppm
Zinc, not less than .....	_____	ppm

1 If added.

2 Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

#### INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials (AAFCO).

### MIXING DIRECTIONS

Thoroughly mix Type B Medicated Swine feed with feed to provide a Type B medicated feed containing up to 36,300 grams tilmicosin per ton or to provide a complete Type C medicated feed containing 181 to 363 grams tilmicosin phosphate per ton. Do not use in any feeds containing bentonite. Bentonite in feeds may affect the efficacy of tilmicosin.

#### Type B Medicated feed to another concentration of Type B Medicated Feed

Starting concentration of Bluebird Type B Medicated Feed		Amount of Type B Medicated Feed to add per ton		Resulting concentration in Type B Medicated Feed	
Grams per ton	Grams per pound	Pounds	Grams per ton	Grams per pound	
36,300	18.1	1,500	27,200	13.6	
		1,000	18,100	9.1	
27,200	13.6	1,330	18,100	9.05	

#### Type B Medicated feed to Type C Medicated feed

Starting concentration of Bluebird Type B Medicated Feed		Amount of Type B Medicated Feed to add per ton	Resulting concentration in Type C Medicated Feed <sup>a</sup>
Grams per ton	Grams per pound	Pounds	Grams per ton
36,300	18.1	20	363
		15	272
		10	181
27,200	13.6	26.7	363
		20	272
		13.3	181
18,100	9.05	40.1	363
		30.1	272
		20	181

**CAUTION:**

Feed containing tilmicosin shall not be fed to pigs for more than 21 days during each phase of production without ceasing administration for re-evaluation of antimicrobial use by a licensed veterinarian before re-initiating a further course of therapy with an appropriate antimicrobial. Veterinary Feed Directive (VFD) expiration date must not exceed ninety (90) days from the time of issuance. VFDs for tilmicosin phosphate shall not be refilled.

Do not allow horses or other equines access to feeds containing tilmicosin. The safety of tilmicosin has not been established in male swine intended for breeding purposes.

**WARNINGS:**

**RESIDUE WARNING:** Swine intended for human consumption must not be slaughtered within 7 days of the last treatment with this drug product.

**Manufactured By  
Bluebird Feed Company  
Any town, USA 12345**

**Net Weight:   lb (  kg)**

For emergency medical information, to report an adverse effect, or for technical service call: 1-800-428-4441.

<sup>a</sup> Sourced from Pulmotil<sup>®</sup>, NADA# 141-064

<sup>\*</sup> Final printed label must bear a single drug concentration.

Elanco, Pulmotil, and the diagonal color bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

