

N-140942-E-0007 (CUM Retyped) Approved 3/8/2001

Narasin and Nicarbazine/Bambermycins Broiler Chicken Ration

Type C Med. Feed

page 1 of 2

Net weight lb (kg) on bag or bulk

Narasin and Nicarbazine/Bambermycins Broiler Chicken Ration Type C Medicated Feed

For the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*, and for increased rate of weight gain and improved feed efficiency in broiler chickens.

Active Drug Ingredient

Narasin.....	27 to 45 g/ton
Nicarbazine.....	27 to 45 g/ton
Bambermycins.....	1.0 to 2.0 g/ton

Guaranteed Analysis

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than.....	_____	%
Sodium ² , not more than.....	_____	%

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

Directions for Use

Feed continuously as sole ration. Narasin and nicarbazine can be combined only at a 1:1 ratio for the 27 to 45 g/ton range.

WARNING: Withdraw five (5) days prior to slaughter.

(OVER)

N-140942-E-0007 (CVM Retyped) Approved 3/8/2001

Narasin and Nicarbazine/Bambermycins Broiler Chicken Ration

Type C Med. Feed

page 2 of 2

CAUTION: For broiler chickens only. Nicarbazine medicated broiler chickens may show reduced heat tolerance when exposed to high temperature and high ambient humidity. Provide adequate drinking water and ventilation during these periods. Do not allow adult turkeys, horses, or other equines access to formulations containing narasin. Ingestion of narasin by these animals has been fatal. Do not feed to laying hens.

MANUFACTURED BY

BLUE BIRD FEED MILL

Anytown, USA 12345