

N-141097-A-0000 (CVM Re typed) Approved 2/3/1999  
Bluebird Swine Type B Med Feed (Ivermectin 364-910 g/ton, BMD 1.0-2.5 g/lb.)

Ivermectin/Bacitracin Methylene Disalicylate Combination  
in Medicated Feed for Swine Type B Medicated Feed

Page 5  
Bluebird Labeling

Page 1 of 2

BLUEBIRD  
SWINE TYPE B MEDICATED FEED

For the treatment and control of gastrointestinal roundworms (*Ascaris suum*, adults and fourth-stage larvae; *Ascarops strongylina*, adults; *Hyostrongylus rubidus*, adults and fourth-stage larvae; *Oesophagostomum* spp., adults and fourth-stage larvae), kidneyworms (*Stephanurus dentatus*, adults and fourth-stage larvae), lungworms (*Metastrongylus* spp., adults), threadworms (*Strongyloides ransomi*, adults and somatic larvae, and prevention of transmission of infective larvae to piglets, via colostrum or milk, when fed during gestation), lice (*Haematopinus suis*) and mange mites (*Sarcoptes scabiei* var. *suis*).

For increased rate of weight gain and improved feed efficiency in growing and finishing swine.

ACTIVE DRUG INGREDIENTS

Ivermectin..... 364-910 g/ton  
Bacitracin methylene disalicylate ..... 1.0-2.5 g/lb

GUARANTEED ANALYSIS

Crude Protein, not less than \_\_\_\_\_%  
Crude Fat, not less than \_\_\_\_\_%  
Crude Fiber, not more than \_\_\_\_\_%

INGREDIENTS

Each ingredient must be specifically named (unless stated as such in the guaranteed analysis listing) in accordance with the names and definitions adopted by AAFCO. Collective terms as listed in 21 CFR 501.110 may be used where applicable.

MIXING DIRECTIONS

This Type B medicated feed must be mixed with feed ingredients (or unmedicated feed) at the rate of 10 to 4 pounds per ton to manufacture a Type C medicated feed containing 1.8 grams of ivermectin and 10 grams of bacitracin methylene disalicylate per ton.

IMPORTANT

Must be thoroughly mixed in feeds before use.  
Store at room temperature.

N-141097-A 0000 (CVM Retyped) Approved 2/3/1999  
Bluebird Swine Type B Med Feed (Ivermectin 364-960 g/ton, BMD 1.0-2.5 g/lb.)  
page 2 of 2

Ivermectin/Bacitracin Methylene Disalicylate Combination  
in Medicated Feed for Swine Type B Medicated Feed

DIRECTIONS FOR USE

Weaned, Growing & Finishing Pigs. Feed the Type C medicated feed, containing bacitracin methylene disalicylate at the recommended level of 10 g/ton and ivermectin at the recommended level of 1.8 g/ton (approximately 100 mcg ivermectin per kg body weight per day) as the only feed for 7 consecutive days. A Type C medicated feed containing bacitracin methylene disalicylate should be fed as the only feed for the duration of the bacitracin methylene disalicylate treatment for increased rate of weight gain and improved feed efficiency in growing and finishing swine from weaning to market weight.

**WARNING:** Withdraw 5 days before slaughter.  
Keep this and all drugs out of the reach of children.

**CAUTION:** Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

This product contains IVOMEC® brand ivermectin and has been formulated specifically for use in swine only. This product should not be used for other animal species. Do not permit water runoff from swine production sites to directly enter lakes, streams, or ponds. Do not contaminate water by direct application or by the improper disposal of drug containers. Dispose of containers in an approved landfill or by incineration.

IVOMEC® is a registered trademark of Merial Limited;  
BMD® is a registered trademark of Alpharma, Inc.

Manufactured by

BLUEBIRD FEED COMPANY  
City, State, Zip Code

Net Weight on bag or bulk