

page 1 of 2

BLUEBIRD  
SWINE TYPE B MEDICATED FEED

For the treatment and control of gastrointestinal roundworms (*Ascaris suum*, adults and fourth-stage larvae; *Ascarops strongylina*, adults; *Hyostrogylus rubidus*, adults and fourth-stage larvae; *Oesophagostomum* spp., adults and fourth-stage larvae), kidneyworms (*Stephanurus dentatus*, adults and fourth-stage larvae), lungworms (*Metastrongylus* spp., adults), threadworms (*Strongyloides ransomi*, adults and somatic larvae, and prevention of transmission of infective larvae to piglets, via colostrum or milk, when fed during gestation), lice (*Haematopinus suis*) and mange mites (*Sarcoptes scabiei* var. *suis*).

For increased rate of weight gain and improved feed efficiency in growing and finishing swine.

ACTIVE DRUG INGREDIENTS

Ivermectin.....	121-728 g/ton
Bacitracin methylene disalicylate .....	1.0-6.0 g/lb

GUARANTEED ANALYSIS

Crude Protein, not less than	___%
Crude Fat, not less than	___%
Crude Fiber, not more than	___%

INGREDIENTS

Each ingredient must be specifically named (unless stated as such in the guaranteed analysis listing) in accordance with the names and definitions adopted by AAFCO. Collective terms as listed in 21 CFR 501.110 may be used where applicable.

MIXING DIRECTIONS

This Type B medicated feed must be mixed with feed ingredients (or unmedicated feed) at the rate of 30 to 5 pounds per ton to manufacture a Type C medicated feed containing 1.8 grams of ivermectin and 30 grams of bacitracin methylene disalicylate per ton.

IMPORTANT

Must be thoroughly mixed in feeds before use.  
Store at room temperature.

### DIRECTIONS FOR USE

Weaned, Growing & Finishing Pigs. Feed the Type C medicated feed, containing bacitracin methylene disalicylate at the recommended level of 30 g/ton and ivermectin at the recommended level of 1.8 g/ton (approximately 100 mcg ivermectin per kg body weight per day) as the only feed for 7 consecutive days. A Type C medicated feed containing bacitracin methylene disalicylate should be fed as the only feed for the duration of the bacitracin methylene disalicylate treatment for increased rate of weight gain and improved feed efficiency in growing and finishing swine from weaning to market weight.

**WARNING:** Withdraw 5 days before slaughter.  
Keep this and all drugs out of the reach of children.

**CAUTION:** Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

This product contains IVOMEC<sup>®</sup> brand ivermectin and has been formulated specifically for use in swine only. This product should not be used for other animal species. Do not permit water runoff from swine production sites to directly enter lakes, streams, or ponds. Do not contaminate water by direct application or by the improper disposal of drug containers. Dispose of containers in an approved landfill or by incineration.

IVOMEC<sup>®</sup> is a registered trademark of Merial Limited;  
BMD<sup>®</sup> is a registered trademark of Alpharma, Inc.

Manufactured by

BLUEBIRD FEED COMPANY  
City, State, Zip Code

Net Weight on bag or bulk