

N-141097-A-0000 (CVM Retyped) Approved: 2/3/1999

Blue Bird Swine Type B Med. Feed (Ivermectin 14.5-364 g/ton) BMD (1.0-2.5g/lb)

Page 1 of 2

Ivermectin/Bacitracin Methylene Disalicylate Combination
in Medicated Feed for Swine Type B Medicated Feed

Swine dysentery
Page 1

Bluebird Labeling

BLUEBIRD
SWINE TYPE B MEDICATED FEED

For the treatment and control of gastrointestinal roundworms (*Ascaris suum*, adults and fourth-stage larvae; *Ascarops strongylina*, adults; *Hyoststrongylus rubidus*, adults and fourth-stage larvae; *Oesophagostomum* spp., adults and fourth-stage larvae), kidneyworms (*Stephanurus dentatus*, adults and fourth-stage larvae), lungworms (*Metastrongylus* spp., adults), threadworms (*Strongyloides ransomi*, adults and somatic larvae, and prevention of transmission of infective larvae to piglets, via colostrum or milk, when fed during gestation), lice (*Haematopinus suis*) and mange mites (*Sarcoptes scabiei* var. *suis*).

For control of swine dysentery associated with *Treponema hyodysenteriae* on premises with a history of swine dysentery but where signs of disease have not yet occurred; or following an approved treatment of the disease condition.

ACTIVE DRUG INGREDIENTS

Ivermectin..... 14.5-364 g/ton
Bacitracin methylene disalicylate 1.0-25 g/lb

GUARANTEED ANALYSIS

Crude Protein, not less than _____ %
Crude Fat, not less than _____ %
Crude Fiber, not more than _____ %

INGREDIENTS

Each ingredient must be specifically named (unless stated as such in the guaranteed analysis listing) in accordance with the names and definitions adopted by AAFCO. Collective terms as listed in 21 CFR 501.110 may be used where applicable.

MIXING DIRECTIONS

This Type B medicated feed must be mixed with feed ingredients (or unmedicated feed) at the rate of 250 to 10 pounds per ton to manufacture a Type C medicated feed containing 1.8 grams of ivermectin and 250 grams of bacitracin methylene disalicylate per ton.

IMPORTANT

Must be thoroughly mixed in feeds before use.
Store at room temperature.

DIRECTIONS FOR USE

Weaned, Growing & Finishing Pigs. Feed the Type C medicated feed, containing bacitracin methylene disalicylate at the recommended level of 250 g/ton and ivermectin at the recommended level of 1.8 g/ton (approximately 100 mcg ivermectin per kg body weight per day) as the only feed for 7 consecutive days. A Type C medicated feed containing bacitracin methylene disalicylate at the recommended level of 250 g/ton should be fed as the only feed for the duration of the bacitracin methylene disalicylate treatment. Diagnosis should be confirmed by a veterinarian when results are not satisfactory.

WARNING: Withdraw 5 days before slaughter.
Keep this and all drugs out of the reach of children.

CAUTION: Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

This product contains IVOMEC® brand ivermectin and has been formulated specifically for use in swine only. This product should not be used for other animal species. Do not permit water runoff from swine production sites to directly enter lakes, streams, or ponds. Do not contaminate water by direct application or by the improper disposal of drug containers. Dispose of containers in an approved landfill or by incineration.

IVOMEC® is a registered trademark of Merial Limited;
BMD® is a registered trademark of Alpharma, Inc.

Manufactured by

BLUEBIRD FEED COMPANY
City, State, Zip Code

Net Weight on bag or bulk