

N-141085-R-0010 Approved 11/30/2001

Blue Bird Zoalene/BMD-SV2C

Type C Replacement Chicken Grower Feed Medicated

Lot No. _____

NET WEIGHT ON BAG OR BULK

**BLUE BIRD ZOALENE/BMD- SV2C
TYPE C REPLACEMENT CHICKEN GROWER FEED
MEDICATED**

Development of active immunity to coccidiosis and as an aid in the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to bacitracin

ACTIVE DRUG INGREDIENTS

Zoalene 75.4 to 113.5 g/ton
Bacitracin methylene disalicylate 100 to 200 g/ton

GUARANTEED ANALYSIS

| | | |
|---|-------|---|
| Crude Protein, not less than | _____ | % |
| Lysine, not less than | _____ | % |
| Methionine, not less than | _____ | % |
| Crude Fat, not less than | _____ | % |
| Crude Fiber, not more than | _____ | % |
| Calcium, not less than | _____ | % |
| Calcium, not more than | _____ | % |
| Phosphorus, not less than | _____ | % |
| Salt ¹ , not less than | _____ | % |
| Salt ¹ , not more than | _____ | % |
| Sodium ² , not less than | _____ | % |
| Sodium ² , not more than | _____ | % |

¹If added

²Shall be guaranteed only when total Sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed continuously as the sole ration where severe exposure to coccidiosis is expected. To control necrotic enteritis, start this feed at first clinical signs of disease; vary bacitracin dosage based on severity of infection; administer continuously for 5-7 days or as long as clinical signs persist, then reduce bacitracin to prevention level (50 g/ton).

CAUTION: Do not feed to birds over 14 weeks of age. Consult a veterinarian or pathologist if losses exceed 0.5% in a two-day period.

Blue Bird Feed Mills
Robin, IN 12345