

N-141085-R-0010 Approved 11/30/2001

Blue Bird Zoalene/BMD-LMC

Type C Replacement Chicken Starter Feed Medicated Lot No. _____

NET WEIGHT ON BAG OR BULK

BLUE BIRD ZOALENE/BMD- LMC
TYPE C REPLACEMENT CHICKEN STARTER FEED
MEDICATED

Development of active immunity to coccidiosis and as an aid in the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to bacitracin

ACTIVE DRUG INGREDIENTS

Zoalene 75.4 to 113.5 g/ton
Bacitracin methylene disalicylate 100 to 200 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than	_____	%
Lysine, not less than	_____	%
Methionine, not less than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than	_____	%
Salt ¹ , not less than	_____	%
Salt ¹ , not more than	_____	%
Sodium ² , not less than	_____	%
Sodium ² , not more than	_____	%

¹If added

²Shall be guaranteed only when total Sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed continuously as the sole ration where light to moderate exposure to coccidiosis is expected. To control necrotic enteritis, start this feed at first clinical signs of disease; vary bacitracin dosage based on severity of infection; administer continuously for 5-7 days or as long as clinical signs persist, then reduce bacitracin to prevention level (50 g/ton).

CAUTION: Consult a veterinarian or pathologist if losses exceed 0.5% in a two-day period.

Blue Bird Feed Mills
Robin, IN 12345