

N-095543-S-0027

Amprolium/Ethopabate/Bambermycins ~~for~~ Broiler Chicken Ration
Type C Med. Feed

15

**Amprolium/Ethopabate/Bambermycins
Broiler Chicken Ration**

Type C Medicated Feed

As an aid in the prevention of coccidiosis where severe exposure to coccidiosis from *Eimeria acervulina*, *E. maxima*, and *E. brunetti* is likely to occur; for increased rate of weight gain, and improved feed efficiency.

ACTIVE DRUG INGREDIENTS

| | |
|-------------------|-------------|
| Amprolium | 113.5 g/ton |
| Ethopabate..... | 36.3 g/ton |
| Bambermycins..... | 1-3 g/ton* |

*The final printed feed label should list only a single drug concentration.

GUARANTEED ANALYSIS

| | |
|---|---|
| Crude Protein, not less than..... | % |
| Lysine, not less than | % |
| Methionine, not less than..... | % |
| Crude fat, not less than | % |
| Crude fiber, not more than..... | % |
| Calcium, not less than..... | % |
| Calcium, not more than | % |
| Phosphorus, not less than | % |
| Salt ¹ , not less than..... | % |
| Salt ¹ , not more than | % |
| Sodium ² , not less than | % |
| Sodium ² , not more than | % |

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials

DIRECTIONS FOR USE

Feed continuously as the sole ration.

WARNING: Use as the sole source of amprolium.

Net Weight lb. (kg) on bag or bulk

Manufactured by
BlueBird Feed Mill
Anytown, Anystate 12345