N-098340-C-0003 Approved 8/16/1985
Blue Bird Broiler Feed Medicated (COCCI/WG/FE)
Type C Med. Feed.
50 LBS. NET WEIGHT

#### BLUE BIRD BROILER FEED MEDICATED

For the prevention of coccidiosis caused by <u>E. tenella</u>, <u>E. necatrix</u>, <u>E. acervulina</u>, <u>E. maxima</u>, <u>E. mivati</u> and <u>E. brunetti</u>; For increased rate of weight gain and for improved feed efficiency.

## **Active Drug Ingredient**

Bambermycins (Flavomycin<sup>R</sup>) 1-2 grams per ton Monensin (Coban) 90-110 grams/ton

### Guaranteed Analysis

Crude Protein, not less than \_\_\_\_\_\_
Crude Fat, not less than \_\_\_\_\_
Crude Fiber, not less than \_\_\_\_\_

## Ingredients

Ground corn
Ground milo
Corn gluten meal
Wheat middlings
Dehydrated alfalfa meal
Fish meal
Animal fat with BHT
Soybean Oil Meal

Riboflavin supplement
Calcium pantothenate
Niacin, Vit. B<sub>12</sub> supplement
Vit. A & D<sub>3</sub>
Ground limestone 1% Salt ½%
Manganese sulfate 0.02%
Defluorinated phosphate ½%
Potassium iodide 0.002%

# Feeding Directions

Feed continuously as the sole ration

WARNING:

Withdraw three (3) days before slaughter. Do not feed to laying chickens. Do not allow horses or other equine access to feeds containing monensin. Ingestion of monensin by equine has been fatal.

Manufactured by BLUE BIRD FEED MILLS Robin, Indiana

