

MONENSIN SODIUM – OXYTETRACYCLINE
 TYPE C
 BLUE BIRD
 CHICKEN FEED B
 MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

As an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati* and *E. maxima*; and for the control of complicated chronic respiratory disease (CRD or air sac infection) caused by *Mycoplasma gallisepticum* and *Escherichia coli* in broiler chickens.

ACTIVE DRUG INGREDIENT

Monensin sodium.....	110 g/ton
Oxytetracycline	200 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min).....	_____ %
Crude Lysine (Min)	_____ %
Methionine (Min)	_____ %
Crude Fat (Min).....	_____ %
Crude Fiber (Min).....	_____ %
Calcium (Min)	_____ %
Calcium (Max).....	_____ %
Phosphorus (Min)	_____ %
Salt, if added (Min).....	_____ %
Salt, if added (Max).....	_____ %
Total Sodium (Min).....	_____ %
Total Sodium (Max)	_____ %

(Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee)

INGREDIENTS

Ingredients as defined by AAFCO

FEEDING DIRECTIONS

Feed continuously as the sole ration.

WARNING: WITHDRAW 72 HOURS BEFORE SLAUGHTER. DO NOT FEED TO LAYING CHICKENS. DO NOT FEED TO CHICKENS OVER 16 WEEKS OF AGE. DO NOT USE IN LOW-CALCIUM FEED CONTAINING LESS THAN 0.55% DIETARY CALCIUM. USE IN SUCH FEEDS MAY RESULT IN VIOLATIVE RESIDUES.

CAUTION: Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. In the absence of coccidiosis in broiler chickens, use of monensin with no withdrawal period may limit feed intake, resulting in reduced weight gain. Not for replacement chickens intended to become broiler breeding chickens.

_____lbs (____kg) NET WEIGHT

BLUE BIRD FEED MILL
 Robin, IN 00000