

OXYTETRACYCLINE
TYPE C
BLUE BIRD
SHEEP FEED
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For sheep for treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENT

Oxytetracycline500 to 1200 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min) %
Equivalent crude protein from Non-Protein Nitrogen (NPN) when added (max) %
Crude Fat (Min) %
Crude Fiber (Max) %
Calcium, if added (Min) %
Calcium, if added (Max) %
Phosphorus, if added (Min) %
Salt, if added (Min) %
Salt, if added (Max) %
Sodium (Min)* %
Sodium (Max)* %
Copper (Min) ppm
Copper (Max) ppm
Selenium, if added (Min) ppm
Vitamin A, if added (Min) IU/lb

* Minimum and maximum percentage of total Sodium shall be guaranteed only when total Sodium exceeds that furnished by the maximum salt guarantee.

**Minimum and maximum copper if added, or if total copper exceeds 20 ppm.

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed 10 mg oxytetracycline per pound of body weight daily continuously for 7 – 14 days. For example, if a lamb weighs 60 lb, consuming 1 lb of supplement per head per day containing 1200 g/ton oxytetracycline.

WARNING: WITHDRAW 5 DAYS BEFORE SLAUGHTER.

____ lbs (____ kg) NET WEIGHT

BLUE BIRD FEED MILL
Robin, IN 00000