

N-039417-5-0086 Approved 9/4/2002

Blue Bird DECCOX #

Type C Complete Sheep + Goat

Feed Medicated.

**Blue Bird Deccox Type C
Complete Sheep & Goat Feed
MEDICATED**

Sheep: For the prevention of coccidiosis in young sheep caused by *Eimeria ovinoidalis*, *E. parva*, *E. bakuensis*, and *E. crandallis*.

Goats: For the prevention of coccidiosis in young goats caused by *Eimeria christensenii* and *E. ninakohlyakimovae*.

ACTIVE DRUG INGREDIENT

Decoquinatate 12.9 to 90.8 g/ton*

GUARANTEED ANALYSIS

| | |
|---|-------------|
| Crude Protein, not less than | _____ % |
| Equivalent Crude Protein from Non-protein Nitrogen (NPN) ¹ , not more than | _____ % |
| Crude Fat, not less than | _____ % |
| Crude Fiber, not more than | _____ % |
| Calcium, not less than | _____ % |
| Calcium, not more than | _____ % |
| Phosphorus, not less than | _____ % |
| Salt ¹ , not less than | _____ % |
| Salt ¹ , not more than | _____ % |
| Sodium ² , not less than | _____ % |
| Sodium ² , not more than | _____ % |
| Copper ³ , not less than..... | _____ ppm |
| Copper ³ , not more than..... | _____ ppm |
| Selenium, not less than..... | _____ ppm |
| Vitamin A ¹ , not less than | _____ IU/lb |

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

³If added, or if total copper exceeds 20 ppm.

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed at a rate of 22.7 mg decoquinatate per 100 lb of bodyweight (0.5 mg/kg) per day. Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard.

WARNING: Do not feed to sheep and goats producing milk for food.

NET WEIGHT ON BAG OR BULK

BLUE BIRD FEED MILLS
Any Town, USA 12345

=====
*Label must show actual concentration of decoquinatate, which will vary with body weight and feed consumption to provide 22.7 mg/100 lb body weight per day. The following table provides examples of feeding rates and amount of drug per ton of feed.

| Animal Wt. (lbs) | Feed Intake (% Body Wt) | Lbs Feed per Head/day | Decoquinatate (grams/ton) |
|---------------------|----------------------------|--------------------------|------------------------------|
| 25 | 0.5 | .125 | 90.8 |
| | 1.5 | .375 | 30.3 |
| | 3.5 | .875 | 13.0 |
| 50 | 0.5 | .25 | 90.8 |
| | 1.5 | .75 | 30.3 |
| | 3.5 | 1.75 | 13.0 |
| 100 | 0.5 | .50 | 90.8 |
| | 1.5 | 1.50 | 30.3 |
| | 3.5 | 3.50 | 13.0 |