

Amprolium and Ethopabate/Bacitracin Methylene Disalicylate Broiler and Replacement Chicken Ration

Type C Medicated Feed

Where immunity to coccidiosis is not desired; to aid in prevention of coccidiosis where severe exposure to coccidiosis from Eimeria acervulina, E. maxima, and E. brunetti is likely to occur; for increased rate of weight gain in broiler chickens raised in floor pens.

ACTIVE DRUG INGREDIENTS

Table with 2 columns: Ingredient Name and Concentration. Rows include Amprolium (113.5 g/ton), Ethopabate (36.3 g/ton), and Bacitracin Methylene Disalicylate (4-50 g/ton*).

GUARANTEED ANALYSIS

Table with 2 columns: Component and Percentage. Rows include Crude Protein, Lysine, Methionine, Crude fat, Crude fiber, Calcium, Phosphorus, and Sodium (both not less than and not more than).

1If added

2Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials

DIRECTIONS FOR USE

Feed as the sole ration from time chickens are placed on litter until past the time when coccidiosis is ordinarily a hazard.

CAUTION: Not for chickens over 16 weeks of age. Do not feed to laying chickens. Not for use as a treatment for outbreaks of coccidiosis.

WARNING: Use as the sole source of amprolium.

*The final finished feed label should list only a single drug concentration.

Net Weight ___ lb. (___ kg) on bag or bulk

Manufactured by BlueBird Feed Mill Anytown, Anystate 12345