

N-131675-6-0032 Approved 5/31/1994  
Fenbendazole Feral Swine  
Type C Med. Feed  
page 1 of 2

**BLUE BIRD LABEL**

**NET WEIGHT - BAG OR BULK**

**LOT NO. \_\_\_\_\_**

**TYPE C MEDICATED FEED**

**FENBENDAZOLE**

**FERAL SWINE**

**INDICATIONS:** For the control and removal of kidney worms (*Stephanurus dentatus*), round worms (*Ascaris suum*), and nodular worms (*Oesophagostomum dentatum*) in feral swine

**DOSAGE:** 3 mg of fenbendazole/kg body weight/day (1.36 mg/lb/day) for three consecutive days.

**WARNING:** Do not use 14 days before or during the hunting season.

**Active Drug Ingredients**

Fenbendazole 90 to 325 g/ton\*.

\*Only a single level of fenbendazole can be indicated on a feed tag. A different feed tag must be provided for each level of fenbendazole manufactured.

**Guaranteed Analysis**

Crude Protein,	Not less than ___ %
Crude Fat,	Not less than ___ %
Crude Fiber,	Not more than ___ %

**Ingredients**

Each ingredient must be specifically named (unless stated as such in the guaranteed analysis listing) in accordance with the names and definitions adopted by AAFCO

N-131675-G-0032 Approved 5/31/1994

Fenbendazole Feral Swine

Type C Med. Feed

page 2 of 2

Feral Swine (continued)

### Feeding Directions

Feed at a rate to provide 3 mg/kg body weight/day (1.36 mg/lb/day) for three consecutive days.

Type C feeds containing Fenbendazole can be fed to adult and young animals either in a mash or pelleted feed. No prior withdrawal of feed or water is necessary.

The following table gives examples of daily feeding rates based on body weights, daily dosage and grams fenbendazole (FBZ) per ton of feed.

<u>Grams FBZ/ton of feed</u>	<u>Daily Dosage (mg/kg/day)</u>	<u>Feed daily for 3 days the following amount of Type C fenbendazole medicated feed based on total body weight of animal group.</u>		
		<u>Example Group Body Weights</u>		
		<u>100 lbs.</u>	<u>500 lbs.</u>	<u>1,000 lbs</u>
136.0	3.0	2.0 lbs.	10.0 lbs.	20.0 lbs.
272.0	3.0	1.0 lb.	5.0 lbs.	10.0 lbs.

Retreatment may be required in six weeks, depending upon conditions of continued exposure to parasites, condition of treated animals and ambient temperatures.

**Safety:** No contraindications for the use of fenbendazole in a zoo environment have been established. Administration to breeding and pregnant ruminants at 2 to 22 times the recommended dose has had no apparent adverse effect.

**MANUFACTURED BY**

**BLUE BIRD FEED MILLS**

**ROBIN, INDIANA**