

N-049934-S-002022
Clopidol/Bacitracin Zinc for Broiler Chickens
Type B Med. Feed

Clopidol/Bacitracin Zinc
For Broiler Chickens

Type B Medicated Feed

For use in manufacturing medicated chicken feed

Indications for Use: To aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, *E. mivati* in broiler chickens; for increased rate of weight gain and improved feed efficiency.

ACTIVE DRUG INGREDIENTS

Clopidol.....	11.35 g/lb
Bacitracin Zinc.....	0.5 - 2.5 g/lb*

GUARANTEED ANALYSIS

Crude Protein, not less than	_____ %
Lysine, not less than	_____ %
Methionine, not less than	_____ %
Crude fat, not less than	_____ %
Crude fiber, not more than	_____ %
Calcium, not less than	_____ %
Calcium, not more than	_____ %
Phosphorus, not less than	_____ %
Salt ¹ , not less than	_____ %
Salt ¹ , not more than	_____ %
Sodium ² , not less than	_____ %
Sodium ² , not more than	_____ %

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials

MIXING DIRECTIONS

Mix 10 lb. of this Type B medicated feed with 1990 lb. of non-medicated feed to make one ton of Type C medicated feed containing 113.5 g/ton clopidol and 5-25 g/ton* bacitracin zinc.

CAUTION: Do not feed to chickens over 16 weeks of age. Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a 2-day period.

*The final printed feed label should list only a single drug concentration.

Net Weight _____ lb (_____ kg)

Manufactured by

Blue bird Feed Mill
Anytown, Anystate 12345