

Lincomycin SWINE MIX Feed
Type B Medicated Feed
(lincomycin Type B medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the treatment and control of swine dysentery and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*. For reduction in the severity of the effects of respiratory disease associated with *Mycoplasma hyopneumoniae*.

ACTIVE DRUG INGREDIENT

Lincomycin^a (as lincomycin hydrochloride agricultural grade).....201 to 40,000 grams/ton*
[0.1005 to 20 grams/lb]*

GUARANTEED ANALYSIS

Crude Protein, not less than	_____	%
Lysine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than	_____	%
Salt ¹ , not less than	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than.....	_____	%
Sodium ² , not more than.....	_____	%
Selenium, not less than.....	_____	ppm
Zinc, not less than	_____	ppm

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be specifically named in accordance with the names and definitions adopted by the AAFCO.

IMPORTANT

Must be thoroughly mixed in feeds before use.

Do not feed undiluted.

Store bulk feed or open bag in a dry place to prevent caking.

Store at room temperature.

MIXING DIRECTIONS

Complete Feeds:

For treatment of swine dysentery and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*: To make complete feed containing 100 grams of lincomycin, add 995-5 pounds of Lincomycin SWINE MIX Type B medicated feed per ton.

For control of swine dysentery and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*: To make complete feed containing 40 grams of lincomycin, add 398-2 pounds of Lincomycin SWINE MIX Type B medicated feed per ton.

For reduction in the severity of the effects of respiratory disease associated with *Mycoplasma hyopneumoniae*: To make complete feed containing 100-200 grams of lincomycin, add 1990-5 pounds of Lincomycin SWINE MIX Type B medicated feed per ton.

DIRECTIONS FOR USE of Type C Medicated Feeds Manufactured from Lincomycin SWINE MIX Medicated (Type B Medicated Feed)

For Treatment of Swine Dysentery and the Control of Porcine Proliferative Enteropathies (Ileitis) Caused by *Lawsonia intracellularis*.

Feed 100 grams of lincomycin per ton of complete feed as the sole ration for three weeks or until signs of disease (watery, mucoid, or bloody stools) disappear.

For Treatment and Control of Swine Dysentery and the Control of Porcine Proliferative Enteropathies (Ileitis) Caused by *Lawsonia intracellularis*.

Feed 100 grams of lincomycin per ton of complete feed as the sole ration for three weeks or until signs of disease (watery, mucoid, or bloody stools) disappear, followed by 40 grams of lincomycin per ton.

For Control of Swine Dysentery and the Control of Porcine Proliferative Enteropathies (Ileitis) Caused by *Lawsonia intracellularis*.

Feed 40 grams of lincomycin per ton of complete feed as the sole ration. For use in animals or on premises with a history of swine dysentery but where symptoms have not yet occurred.

For Reduction in the Severity of the Effects of Respiratory Disease associated with *Mycoplasma hyopneumoniae*

Feed 100-200 grams of lincomycin per ton of complete feed as the sole ration for 21 days.

CAUTION

Occasionally, swine fed lincomycin may within the first two days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within five to eight days without discontinuing the lincomycin treatment. The effects of lincomycin on swine reproductive performance, pregnancy, and lactation have not been determined.

Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

Good Manufacturing Practices Should Be Observed in Preparing Feeds Containing Lincomycin SWINE MIX Medicated Feed (Both Type B and Type C Medicated Feeds).

This includes appropriate clean-out procedures to avoid cross-contamination.



WARNING: No drug withdrawal period is required before slaughter of swine fed lincomycin at approved concentrations (40, 100, or 100-200 grams lincomycin per ton of feed).



To report adverse effects, access medical information, or obtain additional product information, call 1-888-963-8471.

Lot Number (if applicable): _____

Net Weight _____ lb (_____ kg) Bag or Bulk

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

*The final printed feed label should list only a single drug concentration.

^aLincomix[®] is the proprietary name of lincomycin formulation Type A medicated article (NADA 097-505).

Zoetis Inc. [14 September 2017]