

N-114794-C-0002 Approved 6/12/1984

Blue Bird Broiler Ration (COCCI/FE)

282

Type C Med. Feed

8.

Page 1 of 2

Blue Bird Broiler Ration
Medicated

Broiler chickens: as an aid in the prevention of coccidiosis where severe exposure to coccidiosis from E. acervulina, E. maxima, and E. brunetti is likely to occur and for improved feed efficiency.

ACTIVE DRUG INGREDIENTS

Amprolium	0.0125%
Ethopabate	0.004%
Zinc Bacitracin	4-50 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than	18%
Crude Fat, not less than	4%
Crude Fiber, not more than	3%

MANUFACTURER

Blue Bird Feed Mills
Robinwood, Indiana
47907

IMPORTANT

See Reverse Side of Tag for Feeding Directions

N-114794-C-0002 Approved 6/12/1984 ²⁸³
Blue Bird Broiler Ration (COCCI/FE)
Type C Med. Feed
Page 2 of 2

9.

FEEDING DIRECTIONS

Feed as the sole ration from the time
chicks are placed on litter to market weight

If losses exceed 0.5% in a 2-day period, obtain an accurate
diagnosis and follow the instructions of your veterinarian
or poultry pathologist.

WARNING

Do not feed to laying chickens. Use as the
sole source of amprolium.

Do not use this feed as a treatment for outbreaks
of coccidiosis.

cc: Original, NADA
Dup, NADA —
HFA-305
OGC-1
HFV-415 (FOI Staff)
HFV-320
HFV-410 (Matheson)
MCTHOMAS, HFV-128:ljm:4/24/84
revised 5/9/84