

**Tylosin Swine Ileitis Control Medicated Feed  
Type C Medicated Feed**

**CAUTION:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

**INDICATION**

*For control of porcine proliferative enteropathies (ileitis) associated with Lawsonia intracellularis.*

**ACTIVE DRUG INGREDIENT**

Tylosin phosphate <sup>†</sup> ..... 40 or 100 g/ton\*

**GUARANTEED ANALYSIS**

Crude Protein, not less than .....	_____	%
Lysine, not less than .....	_____	%
Crude Fat, not less than .....	_____	%
Crude Fiber, not more than .....	_____	%
Calcium, not less than .....	_____	%
Calcium, not more than .....	_____	%
Phosphorus, not less than .....	_____	%
Salt <sup>1</sup> , not less than .....	_____	%
Salt <sup>1</sup> , not more than .....	_____	%
Sodium <sup>2</sup> , not more than .....	_____	%
Sodium <sup>2</sup> , not less than .....	_____	%
Selenium, not less than .....	_____	ppm
Zinc, not less than .....	_____	ppm

<sup>1</sup> If added.

<sup>2</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**CAUTION**

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

**FEEDING DIRECTIONS**

Feed 100 g of tylosin per ton of complete feed for 21 days. Alternatively, feed 100 g of tylosin per ton of complete feed for at least three weeks, followed by 40 g tylosin per ton of complete feed until pigs reach market weight. Feed continuously as the sole ration.

**MANUFACTURED BY  
BLUE BIRD FEED MILL  
Any town, USA 12345**

**Net Weight lb (kg) on bag or bulk**

‡ Sourced from Tylovet<sup>®</sup>, ANADA 200-484

\* The medicated feed label must state as single drug concentration.

Tylovet is a registered trademark of Hluwapharma EOOD.