

**Clopidol and Chlortetracycline Broiler and Replacement Chicken Supplement
Type B Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD)
drug to use by or on the order of a licensed veterinarian.**

Do not feed undiluted.

Important: Must be thoroughly mixed into feed before use.

INDICATIONS FOR USE

As an aid in prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. mivati* and *E. brunetti* and for control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to chlortetracycline in broiler chickens and replacement chickens intended for use as caged layers.

For use in manufacturing medicated chicken feeds.

ACTIVE DRUG INGREDIENT

Clopidol ^a	11.35 g/lb
Chlortetracycline ^b	10–20 g/lb*

GUARANTEED ANALYSIS

Crude Protein, not less than	_____ %
Lysine, not less than	_____ %
Methionine, not less than	_____ %
Crude Fat, not less than	_____ %
Crude Fiber, not more than	_____ %
Calcium, not less than.....	_____ %
Calcium, not more than	_____ %
Phosphorus, not less than.....	_____ %
Salt ¹ , not less than	_____ %
Salt ¹ , not more than	_____ %
Sodium ² , not less than	_____ %
Sodium ² , not more than	_____ %

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

MIXING DIRECTIONS

Mix the Type B medicated feed with non-medicated ingredients to manufacture one ton of complete Type C medicated chicken feed containing 113.5 grams clopidol and 100 to 200 grams chlortetracycline (CTC) per ton. The following table provides examples of mixing rates and amounts.

Type B concentration g/lb.		Type B per ton of Type C, lbs.	non-medicated feed per ton of Type C, lbs.	Type C Concentrations g/ton	
Clopidol ^a	CTC ^b			CTC	Clopidol
11.35	10	10	1990	100	113.5
11.35	15	10	1990	150	113.5
11.35	20	10	1990	200	113.5

WARNING: Do not feed to chickens over 16 weeks of age. Do not feed to chickens producing eggs for human consumption.

MANUFACTURED BY

BLUE BIRD FEED MILL

Anytown, USA 12345

Net weight ____ lb (____ kg) on bag or bulk

*The final printed label must bear a single drug concentration.

^a Sourced from COYDEN[®] 25 (NADA 034-393).

^b Sourced from CHLORMAX[®] (NADA 046-699), CHLORTET[®] (NADA 048-480), AUREOMYCIN[®] (NADA 048-761), PFICHLOR[®] (NADA 049-287), CLTC[®] (NADA 092-286), and PENNCHLOR[®] (NADA 138-935).

COYDEN is a registered trademark of Huvepharma EOOD

CHLORTET and PENNCHLOR are registered trademarks of Pharmgate Animal Health, LLC

CHLORMAX, PFICHLOR, and AUREOMYCIN are registered trademarks of Zoetis Inc.

CLTC is a registered trademark of Phibro Animal Health Corporation