

OXYTETRACYCLINE
TYPE B
BLUE BIRD
LOBSTER FEED
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS FOR USE
For control of gaffkemia caused by <i>Aerococcus viridans</i> .

ACTIVE DRUG INGREDIENT

Oxytetracycline.....20 g/lb

GUARANTEED ANALYSIS

Crude Protein (Min).....%
Crude Fat (Min).....%
Crude Fiber (Max).....%
Phosphorus (Min).....%

INGREDIENTS

Ingredients as defined by AAFCO.

MIXING DIRECTIONS – Mix 100 lbs. of this Type B with 1900 lbs of feed to manufacture one ton of Type C feed containing 1 g oxytetracycline/lb of feed. Feed the resulting Type C medicated feed as the sole ration for 5 consecutive days.

► WARNING: Withdraw medicated feed 30 days before harvesting lobsters. ◀

FOR USE IN DRY FEEDS ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS.

___lbs(___kg) NET WEIGHT

BLUE BIRD FEED MILL
ROBIN, IN 00000